

Adjacent nesting by Lammergeier *Gypaetus barbatus* and Himalayan Griffon *Gyps himalayensis* on the Tibetan Plateau, China

TODD E. KATZNER, CHIEN HSUN LAI, JAMES D. GARDINER, J. MARC FOGGIN,
DAVID PEARSON and ANDREW T. SMITH

Old world vultures are thought to partition or compete for several types of resources. For example, in Africa, where vulture species diversity is highest, there is evidence that they compete for food (Petrides 1959, König 1983, Mundy *et al.* 1992, Hertel 1994). In Europe, Eurasian Griffon *Gyps fulvus* compete extensively with Lammergeiers *Gypaetus barbatus* for nest sites (Fernández and Donazar 1991, Margalida and Garcia 1999, Bertran and Margalida 2002). Evidence of interspecific aggression at nest sites also has been observed between Eurasian Griffon and Cinereous Vultures *Aegypius monachus* (Blanco *et al.* 1997), between Eurasian Griffon and Egyptian Vultures *Neophron percnopterus* (Pascual and Santiago 1991), and between Cinereous Vulture and Bearded Vultures (Aykurt and Kiraç 2001).

In spite of the potential for competition, patterns in resource availability and population density can sometimes allow species that might otherwise compete to have considerable overlap in resource use (Wiens 1977, Steenhof and Kochert 1988). On the Qinghai-Xizang (Tibetan) plateau, China, we observed an apparently unusual setting where two, and possibly three, large, potentially competing raptors appear to have bred on a single small cliff.

METHODS

These observations were made on a cliff at 4,066 m on the Tibetan plateau in southern Qinghai province, China, near the Yangtze river. The cliff was comprised of three relatively small faces oriented towards the south-west; we estimated their combined length to be c.200 m and their maximum height to be c.100 m. The

cliff faced towards a c.5 km long valley of alpine meadow grasslands dominated by *Kobresia* spp. (Cyperaceae). The valley is bisected by an unpaved road and is a traditional pasture for local Tibetan nomads, although it was not being grazed at the time. Our observations were conducted with 10x40 binoculars on 4 August 2002, from several points 100–200 m from the cliff. Two observers independently estimated distances on the cliff face; when estimates were not identical, a mean value was chosen. To map the site, photographs of the cliff face were scanned, outlines of the cliff face were traced on the image using Adobe Photoshop, and nest sites were then identified by comparison with field notes and sketches.

RESULTS

We observed one fully developed Lammergeier chick on a ledge in a cavity on a small face projecting from the main body of the cliff (Fig.1, Site A). An adult Lammergeier sat 1.5 m above the chick, in another similarly sized, similarly shaped cavity (Fig.1, Site B). These birds were present both in the morning and afternoon of 4 August, and did not move much or fly during our observation periods. A nearly full-grown Himalayan Griffon chick was seen on a 1.5 m long ledge c.15 m above the Lammergeier nest (Fig.1, Site C). The chick moved regularly, sometimes flapping its wings, but never left the nest site. An adult Himalayan Griffon visited the nest for c.2 minutes and regurgitated food for the chick.

The cliff face had other large nests and many potentially suitable nest and roost sites (Fig.1, Sites X and Y). The extensive whitewash near several of these




Figure 1. Map of cliff face. Sites are marked as follows: A. Lammergeier nest and nestling. B. Adult Lammergeier. C. Himalayan Griffon nest and nestling. D. Saker Falcon roost. X. Empty nests. Y. Other roosts

suggested that other vultures regularly used the cliff for roosting and that they may have bred there earlier in the summer. Furthermore, we observed eight adult and one immature Himalayan Griffons fly from this area of the cliff between 10h00 and 11h00. We also found two nests on an adjacent cliff face (80 m from the left edge of the face in Fig.1). Their construction suggested that they had been built by either Upland Buzzard *Buteo hemilasius* or Common Ravens *Corvus corax*. We were unable to determine if these nests had been used in 2002.

In addition we observed a Saker Falcon *Falco cherrug* at the cliff exhibiting apparently territorial behaviour. When we approached the cliff, the bird flew repeatedly from a roost (Fig. 1, Site D), circled above the area calling extensively, and returned to the cliff face. Although we were unable to determine the age of this bird or to locate a nest, this bird's behaviour was consistent with that observed at Saker Falcon nests with chicks in other locations in the region.

DISCUSSION

We found active nests of one Lammergeier and at least one Himalayan Griffon in close proximity, with the strong suggestion that Saker Falcon also bred at this site. Our observations are unusual because nest-based interspecific aggression between vulture species or between vultures and other raptors is frequently strong enough to have negative reproductive consequences (Fernandez and Donazar 1991, Pascual and Santiago 1991, Margalida and Garcia 1999, Aykurt and Kiraç 2001, Matus 2002).

Lammergeiers are solitary breeders, often occupying nests high on cliffs in mountains or river valleys (Snow and Perrins 1998). Their nests are often, but not always, well-spaced from conspecifics and other vultures (Margalida and Garcia 1999, Bertran and Margalida 2002). Himalayan Griffon is one of the world's least-known vultures. Some reports suggest that they are not colonial breeders (Flint *et al.* 1989), while others suggest that they are semi-colonial (Ferguson-Lees and Christie 2001). Saker Falcons are solitary breeders, and although often aggressive towards other raptors in their breeding area, they will sometimes nest near to them (Ellis *et al.* 1997).

All three species probably nest at the same time on the Tibetan plateau. Based on observed fledging dates between 1 and 10 August 2002 at several nests in the general vicinity (TK, unpublished data), and breeding timetables from Europe (Snow and Perrins 1998), we estimate that egg-laying by Lammergeiers on the Tibetan plateau occurs in late February or early March. For Himalayan Griffon, if the duration of the breeding cycle is similar to that of Eurasian Griffons in Europe, then initiation of nesting on the Tibetan plateau should occur in about early March (Snow and Perrins 1998). Saker Falcon chicks have been observed in nests on the plateau in June (E. Potapov *in litt.* 2003) and in nearby lowland central Asia their chicks fledge in early July (TK, unpublished data).

Several characteristics of this region and of these species' biology may interact to allow the nesting behaviour that we observed. First, the Tibetan plateau

is heavily grazed, primarily by domesticated yak *Bos grunniens* and sheep *Ovis* spp., but also by some wild ungulates. These animals probably provide an ample source of carrion for vultures, and facilitate high population densities. Second, although we did not measure the availability of nest sites, we observed few suitable cliffs for nesting sites in this region of the plateau. If nest site availability is low, these birds may have been forced to nest in closer proximity to each other than usual. Third, in nearly all other areas where Lammergeier and Eurasian or Himalayan Griffons co-exist, they share resources with a third species of vulture (Ferguson-Lees and Christie 2001). Because these two species are the only large vultures in this region (Cinereous Vultures are uncommon in this part of Qinghai), interspecific competition may be less than in other places where they co-exist. Finally, Eurasian Griffon colony size and level of aggression by Lammergeiers have been shown to be positively correlated (Bertran and Margalida 2002). Himalayan Griffons, because of their semi- or non-colonial nesting, may provoke less response from Lammergeiers.

ACKNOWLEDGEMENTS

We thank the Biodiversity Working Group of the China Council for International Cooperation on Environment and Development for supporting our travel and the Upper Yangtze Organization for providing logistical support. An anonymous reviewer helped to improve the quality of the manuscript.

REFERENCES

- Aykurt, A. and Kiraç, C. O. (2001) Apparent predation attempt by a Lammergeier *Gypaetus barbatus* on Black Vulture *Aegypius monachus* chick in Turkey. *Sandgrouse* 23: 140.
- Bertran, J. and Margalida, A. (2002) Territorial behavior of Bearded Vultures in response to Griffon Vultures. *J. Field Ornithol.* 73: 86–90.
- Blanco, G., Traverso, J. M., Marchamalo, J. and Martinez, F. (1997) Interspecific and intraspecific aggression among Griffon and Cinereous Vultures at nesting and foraging sites. *J. Raptor Res.* 31: 77–79.
- Ellis D. H., Ellis M. H. and Tsengeng P. (1997) Remarkable Saker Falcon (*Falco cherrug*) breeding records for Mongolia. *J. Raptor Res.* 31: 234–240.
- Fernández, C. and Donazar J. A. (1991) Griffon Vultures *Gyps fulvus* occupying eyries of other cliff-nesting raptors. *Bird Study* 38: 42–44.
- Ferguson-Lees, J. and Christie, D. A. (2001) *Raptors of the world*. Boston: Houghton Mifflin.
- Flint, V. E., Bourso-Leland, N. and Kuznetsov, A. A. (1989) *A field guide to birds of Russia and adjacent territories*. Princeton: Princeton University Press.
- Hertel, F. (1994) Diversity in body size and feeding morphology within past and present vulture assemblages. *Ecology* 75: 1074–1084.
- König, C. (1983) Interspecific and intraspecific competition for food among old world vultures. Pp. 153–171 in S. R. Wilbur and J. A. Jackson, eds. *Vulture biology and management*. Berkeley: University of California Press.
- Margalida, A. and Garcia, D. (1999) Nest use, interspecific relationships and competition for nests in the Bearded Vulture *Gypaetus barbatus* in the Pyrenees: influence on breeding success. *Bird Study* 46: 224–229.
- Matus, A. A. (2002) On the breeding biology of the Griffon Vulture in the Crimea. *Berkut* 11: 121–123.

- Mundy, P., Butchart, D., Ledger, J. and Piper, S. (1992) *The vultures of Africa*. London: Academic Press.
- Pascual, J. and Santiago, J. M. (1991) Egyptian Vultures steal food from nestling Griffon Vultures. *J. Raptor Res.* 25: 96–97.
- Petrides, G. A. (1959) Competition for food between five species of East African vultures. *Auk* 76: 104–106.
- Snow, D. W. and Perrins, C. M. (1998) *The birds of the western Palearctic*. Concise Edition. New York: Oxford University Press.
- Todd E. Katzner, School of Life Sciences, Arizona State University, PO Box 874501, Tempe, AZ, 85287-4501, U.S.A. Present address: Renewable Resources Assessment Group, Department of Environmental Science and Technology, South Kensington Campus, Exhibition Road, Imperial College London, SW7 2AZ, U.K. Email: tkatzner@imperial.ac.uk
- Chien Hsun Lai, David Pearson, and Andrew T. Smith, School of Life Sciences, Arizona State University, PO Box 874501, Tempe, AZ, 85287-4501, U.S.A.
- James D. Gardiner and J. Marc Foggin, Plateau Perspectives, 200 Walnut Avenue, St Lambert, Quebec, J4P 2T1, Canada.
- Steenhof, K. and Kochert, M.N. (1988) Dietary responses of three raptor species to changing prey densities in a natural environment. *J. Animal Ecol.* 57: 37–48.
- Wiens, J. A. (1977) Competition and variable environments. *Amer. Sci.* 65: 590–597.

Hybridisation between Slender-billed Gull *Larus genei* and Black-headed Gull *L. ridibundus* in Irkutsk, Russia

IGOR V. FEFELOV

Slender-billed Gull *Larus genei* breeds from the Iberian peninsula to Central Asia, between 20°N and 50°N. It breeds as far east as western Pakistan, Tajikistan and the Novosibirsk region in western Siberia (Zubakin 1988, Ryabitshev 2001). The species was first recorded around Lake Baikal in June 1989 when an adult male was collected in a colony of Black-headed Gulls in the Selenga delta, in south-western Baikal on 15 June (Tupitsyn and Fefelov 1995). Another was seen in June 1991 between Irkutsk and Angarsk, in a flock of non-breeding individuals of Yellow-legged Gull *L. cachinnans* (Popov and Salovarov 2000). A third record was documented in June 2001 at Gusinoe Ozero lake in Buryatia, south-east of Baikal (Tebb and Ranner 2002). Other vagrants have occurred in east Asia (Moore 2002, Tebb and Ranner 2002).

OBSERVATIONS

On 25 May 2003, we found a Slender-billed Gull near Black-headed Gull colonies in a wetland on the edge of Irkutsk (52°19'N 104°13'E). Black-headed Gulls first bred here during the 1980s, and c.300 pairs have nested since the early 1990s. By the end of the month the Slender-billed Gull was found to be paired to a Black-headed Gull, and building a nest in a small colony of seven pairs of Black-headed Gulls in a marsh with reedmace *Typha latifolia*. On 3 June this contained three eggs measuring 53.4–36.7 mm, 53.0–35.5 mm, and 52.2–35.4 mm. They were brownish-green with blackish spotting. The egg-coloration was at the lighter end of the range found in Black-headed Gull. We assume therefore that the female of the pair was the Black-headed Gull, as the eggs of Slender-billed Gull have a significantly lighter background colour: white with a pinkish or cream, but not green, tinge (Zubakin

1988). The structure and size of the nest showed no consistent differences from nests of Black-headed Gulls in the vicinity.

No displays between the pair were seen. Both birds incubated the eggs. When the Slender-billed Gull was flushed from the nest, it flew 30–50 m away, showed no aggressive behaviour, and called less frequently than Black-headed Gulls normally do. Only twice during our observations did we hear it give the species's distinctive 'ka-ka-ka' alarm call.

On 11 June, the Slender-billed Gull was found dead near the colony, having been shot. It was an adult male in breeding plumage. The following biometrics were taken: wing length: 315 mm, tail length 116 mm, tarsus: 52 mm, bill length: 42 mm. The dimensions of the testicles were 15x8 mm (left) and 8x7 mm (right). The skin has been transferred to the Zoological Institute, St. Petersburg, Russia.

As incubation by single parents is atypical in gulls, the clutch was exchanged with that of a pair of Black-headed Gulls nearby, in order to increase the chance of survival. Two chicks of the mixed pair hatched on 23 June and the third followed on 24 June. In the first few days, their coloration was generally similar to, but lighter than, similarly aged Black-headed Gull chicks. The distinctive grey colour on the upperparts of Slender-billed Gull chicks (Zubakin 1988) was absent. The second chick had less black spotting, and a large (10x12 mm) black patch on the front of the crown instead of many small spots as found on Black-headed Gull chicks. Zubakin (1988) studied Black-headed Gull around Moscow and Slender-billed Gull in the Black Sea, and noted that some Slender-billed Gull chicks may have an almost entirely black head, but that this is never observed in Black-headed chicks. Photographs of the chicks show that they are intermediate in appearance between the two species (V. A.

Zubakin *in litt.* 2003). There appear to be few published photographs of young Slender-billed chicks; one in Sauer (1982) closely resembles the second chick described above.

On 26 June, the two younger chicks were ringed in the nests (the older one had already disappeared). As juveniles, the birds had lighter-coloured upperparts, being greyish rather than brownish as in Black-headed Gulls. The base of the bill was pink contrasting with the dark tip, rather than the greyish-pink bill base and less contrasting dark tip of juvenile Black-headed Gull. The last date when the juveniles were seen was 27 July, when they could fly well.

This appears to be the first case of hybridisation between Slender-billed Gull and Black-headed Gulls, and occurred c.2,000 km east of known breeding areas for Slender-billed Gull.

ACKNOWLEDGEMENTS

Thanks to: Dr I. I. Tupitsyn (Irkutsk State Pedagogical University) for much help in the field and for information; Dr V. A. Zubakin (Institute of Problems of Ecology and Evolution, Moscow) for infor-

mation on Slender-billed Gull; Dr V. M. Loskot (Zoological Institute, Sankt-Peterburg) and G. Tebb (BirdLife Austria) for reviewing the literature; and N. Moores and Kim Hyun-tae for valuable information.

REFERENCES

- Moores, N. (2002) Additions to the South Korean List: observations of species not listed by Lee, Koo and Park (2000). Downloaded from <http://www.wbkenglish.com/newbirds27.asp> on 30 November 2002.
- Popov, V. V. and Salovarov, V. O. (2000) [‘Rare bird species of Angarsk District (Southern Baikal area).’] Pp. 191–194 in Dorzhiev, Ts. Z., ed. *The ornithological observation in Russia. Issue 2.* Ulan-Ude: Buryat University Press. (In Russian.)
- Ryabitsev, V. K. (2001) [‘Birds of Ural, Near-Ural area and Western Siberia.’] Ekaterinburg: Ural University. (In Russian.)
- Sauer, F. (1982) *Vögel 2: Wasservögel.* München: Mosaik Verlag.
- Tebb, G. and Ranner, A. (2002) New and significant bird records from Buryatia, Russia. *Forktail* 18: 101–105.
- Tupitsyn, I. I. and Fefelov, I. V. (1995) [‘New bird species at the Baikal Lake.’] *Ornithologia* 26: 197–198. (In Russian.)
- Zubakin, V. A. (1988) [‘Slender-billed Gull.’] Pp. 105–115 in Il’itchyov, V. D. and Zubakin, V. A., eds. *Birds of USSR: Laridae.* Moscow: Nauka. (In Russian.)

Igor Fefelov, Research Institute of Biology, Irkutsk State University, PO Box 24, 664003, Irkutsk, Russia.
Email: u000438@ic.isu.ru

Notes on the population density and feeding ecology of the Collared Falconet *Microhierax caerulescens* in Buxa Tiger Reserve, West Bengal, India

S. SIVAKUMAR, HILLALJYOTI SINGHA and VIBHU PRAKASH

The Collared Falconet *Microhierax caerulescens* is found from India, Nepal and Bhutan through to Myanmar, Thailand, Laos, Cambodia and Vietnam (del Hoyo *et al.* 1994). In India, the species is found throughout the lower Himalayan foothills from Garhwal eastwards to Assam, the north-eastern hill states and Arunachal Pradesh, usually up to 900 m occasionally to 2,000 m (Ali and Ripley 1987). It occurs in deciduous, moist-deciduous and evergreen forest, and is most often observed in man-made clearings, natural open spaces and forest margins (Ali and Ripley 1987). Relatively little has been published on the species, apart from observations of allopreening (Sparks 1965), and breeding (Naoroji 1997, Kemp and Van Zyl 1998). Here we present some observations on the feeding ecology and population density of the species in Buxa Tiger Reserve, West Bengal, India.

STUDY AREA

Buxa Tiger Reserve (26°30–55’N 89°20–55’E) is located in the north-eastern corner of Jalpaiguri

district, West Bengal, India at 60–1,750m. It covers an area of 760 km², with a core area of 385 km² and a buffer zone of 375 km². The habitat is mainly tropical moist-deciduous forest (Champion and Seth 1968) dominated by sal *Shorea robusta*, plus evergreen forest, riverine forest, scrub, grassland, and plantations of sal, teak *Tectona grandis* and jarul *Lagerstroemia reginae*. The temperature ranges from 12°C to 32°C, and the average annual rainfall is 4,100 mm.

METHODS

Falconets were observed during a study on raptor ecology carried out in the reserve during 1998–2000. A 45 km transect was driven every two weeks from January 2000 to December 2000 from Damanpur to Checko via Panijhora, Pambubusti, Rajabathkawa, Santrabari, Jainty and ‘23rd mile tower’. A 50 m band on each side of the transect was searched, with one person observing each side from a slow-moving (20 km/h) jeep. Each census was carried out during 07h00–11h00. Cloudy and rainy days were avoided.