

## References

- Adams, D.B.** (1982). Time of onset and target of immune reactions in sheep with acquired immunity against *Haemonchus contortus*. *International Journal for Parasitology* **12**: 439-443.
- Adams, D.B., Anderson, B.H. and Windon, R.G.** (1989). Cross-immunity between *Haemonchus contortus* and *Trichostrongylus colubriformis* in sheep. *International Journal for Parasitology* **19**: 717-722.
- Akhmetov, A.A.** (1973a). The helminth fauna of sheep in Kokchetav oblast. *Transactions of the Kazakhstan Scientific Research Veterinary Institute* **15**: 348-350. [In Russian.]
- Akhmetov, A.A.** (1973b). Seasonal and age dynamics of helminthoses of sheep in Kokchetav oblast. *Transactions of the Kazakhstan Scientific Research Veterinary Institute* **15**: 351-354. [In Russian.]
- Akhmetov, A.A.** (1975). Generalities on the helminth fauna of sheep in Kokchetav oblast in comparison with neighbouring and southern oblasts of Kazakhstan. Pp. 118-120 in: *Diseases of Sheep and Means of Their Control in Kazakhstan*. Kazakhstan Scientific Research Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]
- Akhmetov, A.A.** (1982). Helminths of sheep in Kokchetav oblast. Pp. 3-12 in: *Problems of Veterinary Parasitology in Kazakhstan. A Collection of Scientific Transactions*. Kazakhstan Scientific Investigation Veterinary Institute, Almaty. [In Russian.]
- Al Saqur, I., Bairden, K., Armour, J. and Gettinby, G.** (1982). Population study of bovine *Ostertagia* spp. infective larvae on herbage and in soil. *Research in Veterinary Science* **32**: 332-337.
- Albon, S.D., Stien, A., Irvine, R.J., Langvatn, R., Ropstad, E. and Halvorsen, O.** (2002). The role of parasites in the dynamics of a reindeer population. *Proceedings of the Royal Society of London Series B* **269**: 1625-1632.
- Alcaino, H.A. and Baker, N.F.** (1974). Comparison of two flotation methods for detection of parasite eggs in feces. *Journal of the American Veterinary Medical Association* **164**: 620-622.
- Alexander, K.A. and Appel, M.J.G.** (1994). African wild dogs (*Lycaon pictus*) endangered by a canine distemper epizootic among domestic dogs near the Masai-Mara national reserve, Kenya. *Journal of Wildlife Diseases* **30**: 481-485.
- Altaev, A.Kh.** (1953). Studies of the helminth fauna of sheep and goats in Dagestan and observations on the biology of *Trichostrongylus skrjabini*. PhD Thesis, Moscow. [In Russian.]
- Anderson, R.C.** (2000). *Nematode Parasites of Vertebrates: Their Development and Transmission* (2<sup>nd</sup> ed.). CAB International, Oxon, UK. Pp. 96-128.
- Anderson, R.M. and Gordon, D.M.** (1982). Processes influencing the distribution of parasite numbers within host populations with special emphasis on parasite-induced host mortalities. *Parasitology* **82**: 373-398.
- Anderson, R.M. and May, R.M.** (1978). Regulation and stability of host-parasite population interactions. 1. Regulatory processes. *Journal of Animal Ecology* **47**: 219-47.
- Anderson, R.M. and May, R.M.** (1982). *Population Biology of Infectious Diseases*. Springer-Verlag, Berlin.

- Anderson, R.M and May, R.M.** (1991). *Infectious Diseases of Humans: Dynamics and Control*. Oxford University Press, Oxford.
- Anderson, T.J.C., Blouin, M.S. and Beech, R.N.** (1998). Population biology of parasitic nematodes: applications of genetic markers. *Advances in Parasitology* **41**: 219-283.
- Andreeva, N.K.** (1957). *Atlas of Helminths (Strongylates) of Domestic and Wild Ruminants of Kazakhstan*. Veterinary Institute, Kazakhstan Branch of the All-Union Academy of Agricultural Science named after V.I. Lenin, Tashkent, USSR. [In Russian.]
- Armour, J.** (1970). Bovine ostertagiasis: A review. *The Veterinary Record* **86**: 184-190.
- Armour, J.** (1980). The epidemiology of helminth disease in farm animals. *Veterinary Parasitology* **6**: 7-46.
- Armour, J.** (1989). The influence of host immunity on the epidemiology of trichostrongyle infections in cattle. *Veterinary Parasitology* **32**: 5-19.
- Arneberg, P. and Folstad, I.** (1999). Predicting effects of naturally acquired abomasal nematode infections on growth rate and food intake in reindeer using serum pepsinogen levels. *Journal of Parasitology* **85**: 367-369.
- Arneberg, P., Folstad, I. And Karter, A.J.** (1996). Gastrointestinal nematodes depress food intake in naturally infected reindeer. *Parasitology* **112**: 213-219.
- Atabaev, A.Kh.** (1953). Studies on the helminth fauna of sheep and goats in Dagestan and observations on the biology of *Trichostrongylus skrjabini*. PhD Thesis, Moscow. [In Russian.]
- Atabaev, A.O.** (1975). Attempts at the control of haemonchosis of sheep in Turkestan raion of Chimkent oblast. Pp. 116-117 in: *Diseases of Sheep and Means of Their Control in Kazakhstan*. Kazakhstan Scientific Research Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]
- Aumont, G. and Gruner, L.** (1989). Population evolution of the free-living stage of goat gastrointestinal nematodes on herbage under tropical conditions in Guadeloupe (French West Indies). *International Journal for Parasitology* **19**: 539-546.
- Aumont, G., Coulaud, G., Grude, A. and Gruner, L.** (1989). Pasture populations of cattle nematode larvae in Guadeloupe (French West Indies). *International Journal for Parasitology* **19**: 547-554.
- Baker, N.F. and Fisk, R.A.** (1976). Phenotypic differences in female *Ostertagia circumcincta* as demonstrated by isozymes hydrolysing alpha-naphthyl acetate. *American Journal of Veterinary Research* **37**: 323-327.
- Baker, N.F., Fisk, R.A. and Rimbey, C.W.** (1984). Seasonal occurrence of infective nematode larvae in California High Sierra pastures grazed by cattle. *American Journal of Veterinary Research* **45**: 1393-1397.
- Balic, A., Bowles, V.M. and Meeusen, E.N.T.** (2000). The immunobiology of gastrointestinal nematode infections in ruminants. *Advances in Parasitology* **45**: 181-241.
- Bang, K.S., Familton, A.S. and Sykes, A.R.** (1990). Effect of ostertagiasis on copper status in sheep: a study involving use of copper oxide wire particles. *Research in Veterinary Science* **49**: 306-314.

- Banks, D.J.D., Singh, R., Barger, I.A., Pratap, B. and Le Jambre, L.F.** (1990). Development and survival of infective larvae of *Haemonchus contortus* and *Trichostrongylus colubriformis* on pasture in a tropical environment. *International Journal for Parasitology* **20**: 155-160.
- Bannikov, A.G., Zhirnov, L.V., Lebedeva, L.S. and Fandeev, A.A.** (1961). *The Biology of the Saiga*. Moscow. [In Russian.]
- Barger, I.A.** (1985). The statistical distribution of trichostrongylid nematodes in grazing lambs. *International Journal for Parasitology* **15**: 645-649.
- Barger, I.A.** (1989). Genetic resistance of hosts and its influence on epidemiology. *Veterinary Parasitology* **32**: 21-35.
- Barger, I.A.** (1997a). Control by management. *Veterinary Parasitology* **72**: 493-506.
- Barger, I.A.** (1997b). Models as a guide to sustainable worm control. In: *Sustainable Control of Internal Parasites in Ruminants. Animal Industries Workshop, June 1997, Lincoln University (New Zealand)* (Ed. Barrell, G.K.), pp. 203-213.
- Barger, I.A.** (1999). The role of epidemiological knowledge and grazing management for helminth control in small ruminants. *International Journal for Parasitology* **29**: 41-47.
- Barger, I.A. and Le Jambre, L.F.** (1988). Regulation of *Haemonchus contortus* populations in sheep: mortality of established worms. *International Journal for Parasitology* **18**: 269-273.
- Barger, I.A. and Southcott, W.H.** (1975). Control of nematode parasites by grazing management. I. Decontamination of cattle pastures by grazing with sheep. *International Journal for Parasitology* **5**: 39-44.
- Barger, I.A., Le Jambre, L.F., Georgi, J.R. and Davies, H.I.** (1985). Regulation of *Haemonchus contortus* populations in sheep exposed to continuous infection. *International Journal for Parasitology* **15**: 529-533.
- Barger, I.A., Siale, K., Banks, D.J.D. and Le Jambre, L.F.** (1994). Rotational grazing for control of gastrointestinal nematodes of goats in a wet tropical environment. *Veterinary Parasitology* **53**: 109-116.
- Barnard, C.J.** (1990). Pathology and host behaviour. Pp. 34-63 in: *Parasitism and Host Behaviour*. Taylor and Francis, London.
- Barnes, E.H, Dobson, R.J. and Barger, I.A.** (1995). Worm control and anthelmintic resistance: adventures with a model. *Parasitology Today* **2**: 56-63.
- Baryshkinov, G.F., Dmitrieva, E.L., Krakhmalnaya, T.V. and Sher, A.V.** (1998). Origin, evolution and systematics. In: (Eds. Sokolov, V.E. and Zhirnov, L.V.). *The Saiga Antelope. Phylogeny, Systematics, Conservation and Land Use*. Moscow. Pp. 9-24. [In Russian]
- Bates, K.M., Hansen, L.P., Green, S.P., Wallace, D.H. and Green, T.J.** (2000). Potential for cross-transmission of *Dictyocaulus viviparus* between cattle and white-tailed deer. *Journal of Wildlife Diseases* **36**: 774-778.
- Begon, M. and Bowers, R.G.** (1995). Beyond host-pathogen dynamics. In: *Ecology of Infectious Diseases in Natural Populations* (Eds. Grenfell, B.T. and Dobson, A.P.). Cambridge, Cambridge University Press. Pp. 478-509.

- Begon, M., Harper, J.L and Townsend, J.R.** (1996a). *Ecology. Individuals, Populations and Communities*. (3<sup>rd</sup> edn). Blackwell Scientific Publications, Oxford.
- Begon, M., Mortimer, M. and Thompson, D.J.** (1996b). *Population Ecology: a unified study of animals and plants* (3<sup>rd</sup> ed.). Blackwell, Oxford.
- Behnke, J.M.** (1987). Evasion of immunity by nematode parasites causing chronic infections. *Advances in Parasitology* **26**: 1-71.
- Behnke, J.M., Barnard, C.J. and Wakelin, D.** (1992). Understanding chronic nematode infections: evolutionary considerations, current hypotheses and the way forward. *International Journal for Parasitology* **22**: 861-907.
- Bekele, T.** (2002). Epidemiological studies on gastrointestinal helminths of dromedary (*Camelus dromedaries*) in semi-arid lands of eastern Ethiopia. *Veterinary Parasitology* **105**: 139-152.
- Bekenov, A.B., Grachev, Iu.A. and Milner-Gulland, E.J.** (1998). The ecology and management of the Saiga antelope in Kazakhstan. *Mammal Review* **28**: 1-52.
- Belem, A.M.G., Couvillion, C.E., Siefker, C. and Griffin, R.N.** (1993). Evidence for arrested development of abomasal nematodes in white-tailed deer. *Journal of Wildlife Diseases* **29**: 261-265.
- Bellows, T.S.** (1981). The descriptive properties of some models for density dependence. *Journal of Animal Ecology* **50**: 139-156.
- Benton, T.G. and Grant, A.** (1999). Elasticity analysis as an important tool in evolutionary and population ecology. *Trends in Ecology and Evolution* **14**: 467-471.
- Berding C., Keymer AE., Murray JD. and Slater AF.** (1986). The population dynamics of acquired immunity to helminth infection. *Journal of Theoretical Biology*. **122**: 459-471.
- Berding, C., Keymer, A.E., Murray, J.D. and Slater, A.F.G.** (1987). The population dynamics of acquired immunity to helminth infection: experimental and natural transmission. *Journal of Theoretical Biology* **126**: 167-182.
- Berezovski, V.S.** (1975). Seasonal and age dynamics of marshallagiosis of sheep in Kustanai oblast. In: *Diseases of Sheep and Means of Their Control in Kazakhstan*. Kazakhstan Scientific Research Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. Pp. 121-122. [In Russian.]
- Berkinbaev, O.** (1992). Studies on diseases of saigas in Kazakhstan. In *Diseases and Parasites of Wild Animals*. Ministry of Ecology and Natural Resources, All-Russian Scientific Research Institute of Natural Reserves and Reserve Affairs, Centre for the Health of Wild Animals, Moscow. Pp. 32-41. [In Russian.]
- Berkinbaev, O., Baitursunov, K.K., Pinaeva, L.M., Priadko, E.I., Grachev, Y.A., Bekenov, A.B., Izkenov, N.T., Bisenova, P., Tastanov, T.B., Dimkova, N.D. and Kokhno, L.I.** (1994). An ecologo-faunal analysis of endoparasites present in the saiga antelope in Kazakhstan. *Deposition to the Kazakhstan State Institute of Scientific Technical Information*, no. 4937-Ka.94. Almaty, Kazakhstan, 56 pp. [In Russian.]
- Besier, R.B. and Dunsmore, J.D.** (1993). The ecology of *Haemonchus contortus* in a winter rainfall climate in Australia: the survival of infective larvae on pasture. *Veterinary Parasitology* **45**: 293-306.

- Beveridge, I. And Ford, G.E.** (1982). The trichostrongyloid parasites of sheep in South Australia and their regional distribution. *Australian Veterinary Journal* **59**: 177-179.
- Bliss, C.I. and Fisher, R.A.** (1953). Fitting the negative binomial distribution to biological data. *Biometrics* **9**: 176-200.
- Blouin, M.S., Yowell, C.A., Courtney, C.H. and Dame, J.B.** (1995). Host movement and the genetic structure of populations of parasitic nematodes. *Genetics* **141**: 1007-1014.
- Blouin, M.S., Yowell, C.A., Courtney, C.H. and Dame, J.B.** (1997). *Haemonchus placei* and *Haemonchus contortus* are distinct species based on mtDNA evidence. *International Journal for Parasitology* **11**: 1383-1387.
- Boag, B. and Kolb, H.H.** (1989). Influence of host age and sex on nematode populations in the wild rabbit (*Oryctolagus cuniculus* L.). *Proceedings of the Helminthological Society of Washington* **56**: 116-119.
- Boag, B. and Thomas, R.J.** (1975). Epidemiological studies on *Nematodirus* species in sheep. *Research in Veterinary Science* **19**: 263-268.
- Boag, B. and Thomas, R.J.** (1977). Epidemiological studies on gastrointestinal parasites of sheep: the seasonal number of generations and succession of species. *Research in Veterinary Science* **22**: 62-67.
- Boag, B.B., Topham, P.B. and Webster, R.** (1989). Spatial distribution on pasture of infective larvae of the gastrointestinal nematode parasites of sheep. *International Journal for Parasitology* **19**: 681-685.
- Boag, B., Lello, J., Fenton, A., Tompkins, D.M. and Hudson, P.J.** (2001). Patterns of parasite aggregation in the wild European rabbit (*Oryctolagus cuniculus*). *International Journal for Parasitology* **31**: 1421-1428.
- Boev, S.N., Sokolova, I.B. and Panin, V.Y.** (1962). *Helminths of Conserved Ruminants of Kazakhstan*. National Academy of Sciences of Kazakhstan, Almaty, USSR. 377 pp. [In Russian]
- Booth, M., Bundy, D.A.P., Albonico, M., Chwya, H.M., Alawi, K.S. and Savioli, L.** (1998). Associations among multiple geohelminth species infections in schoolchildren from Pemba island. *Parasitology* **116**: 85-93.
- Borgsteede, F.H.M.** (1996). The effect of parasites on wildlife. *The Veterinary Quarterly* **18**: S138-S140.
- Brooker, S. and Michael, E.** (2000). The potential of Geographical Information Systems and remote sensing in the epidemiology of human helminth infections. *Advances in Parasitology* **47**: 246-288.
- Brown, N.J.** (1996). *The New Penguin Russian Course*. Penguin, London. p.11.
- Brundson, R.V.** (1963). The seasonal availability to grazing sheep of infective trichostrongyle larvae on pasture. *New Zealand Veterinary Journal* **11**: 86-89.
- Brundson, R.V.** (1970). Within-flock variations in strongyle worm infections in sheep: the need for adequate diagnostic samples. *New Zealand Veterinary Journal* **18**: 185-188.

- Brundson, R.V.** (1971). Trichostrongyle worm infection in cattle: further studies on problems of diagnosis and on seasonal patterns of occurrence. *New Zealand Veterinary Journal* **19**: 203-212.
- Bryan, R.P. and Kerr, J.D.** (1988). The grazing behaviour of cattle in relation to the sampling of infective nematode larvae on pasture. *Veterinary Parasitology* **30**: 73-82.
- Bryan, R.P. and Kerr, J.D.** (1989a). The relation between the natural worm burden of steers and the faecal egg count differentiated to species. *Veterinary Parasitology* **30**: 327-334.
- Bryan, R.P. and Kerr, J.D.** (1989b). Factors affecting the survival and migration of the free-living stages of gastrointestinal nematode parasites of cattle in Central Queensland. *Veterinary Parasitology* **30**: 315-326.
- Burrows, R.O., Best, P.J. and Preston, J.M.** (1980). Trichostrongylid egg output of dairy cows. *Veterinary Record* **107**: 399-401.
- Bush, A.O., Fernandez, J.C., Esch, G.W. and Seed, J.R.** (2001). *Parasitism : the Diversity and Ecology of Animal Parasites*. Cambridge University Press, Cambridge.
- Bye, K. and Halvorsen, O.** (1983). Abomasal nematodes of the Svalbard reindeer (*Rangifer tarandus platyrhynchus* Vrolik). *Journal of Wildlife Diseases* **19**: 101-105.
- Cabaret, J.** (1984). Seasonal changes in the abomasal nematodes of naturally infected ewes in Moulay-Bouazza (Morocco). *Veterinary Parasitology* **15**: 47-56.
- Case, T.J.** (2000). *An Illustrated Guide to Theoretical Ecology*. Oxford University Press.
- Cerioni, S. and Mariniello, L.** (1996). Parasitic metazoans of *Stenella coeruleoalba* (Cetacea: Delphinidae) stranded along the coast of Latium, 1985-91. *Parassitologia* **38**: 505-10 (Abstract.)
- Chaloupka, M.** (2002). Stochastic simulation modelling of southern Great Barrier Reef green turtle population dynamics. *Ecological Modelling* **148**: 79-109.
- Cherikaeva, N.S. and Martinov, V.V.** (1975). The distribution of helminths of sheep in Ural oblast. Pp. 321-324 in: *Diseases of Sheep and Means of Their Control in Kazakhstan*. Kazakhstan Scientific Research Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]
- Chiejina, S.N. and Emehelu, C.O.** (1984). Seasonal changes in pasture populations of infective larvae of gastrointestinal nematodes of cattle in eastern Nigeria. *Research in Veterinary Science* **37**: 144-147.
- Chiejina, S.N. and Fakae, B.B.** (1989). The ecology of infective larvae of bovine gastrointestinal trichostrongylids in dry season contaminated pastures in the Nigerian derived savanna. *Journal of Helminthology* **63**: 127-139.
- Chilonda, P., Woodford, J.D., Ahmadu, B., Samui, K.L., Syakalima, M. and Mlangwa, J.E.** (1999). Foot and mouth disease in Zambia: a review of the aetiology and epidemiology and recommendations for possible control. *Revue Scientifique et Technique de L'Office International des Epizooties* **18**: 585-592.
- Chitty D.** (1960). Population processes in the vole and their relevance to general theory. *Canadian Journal of Zoology* **38**: 99-113.

- Ciordia, H. and Bizzell, W.E.** (1963). The effects of various constant temperatures on the development of the free living stages of some nematode parasites of cattle. *The Journal of Parasitology* **49**: 60-63.
- Ciordia, H., Neville, W.E., Baird, D.M. and McCampbell, H.C.** (1971). Internal parasitism of beef cattle on winter pastures: level of parasitism as affected by stocking rates. *American Journal of Veterinary Research* **32**: 1353-1358.
- Clark, I.A.** (2001). Heterologous immunity revisited. *Parasitology* **122**: S51-S59.
- Cleaveland, S., Hess, G.R., Dobson, A.P., Laurenson, M.K., McCallum, H.I., Roberts, M.G. and Woodroffe, R.** (2002). The role of pathogens in biological conservation. In: Hudson, P.J., Rizzoli, A., Grenfell, B.T., Heesterbeek, H. and Dobson, A.P. (Eds.). *The Ecology of Wildlife Diseases*. Oxford University Press. Pp. 139-150.
- Coltman, D.W., Pilkington, J., Kruuk, L.E.B., Wilson, K. and Pemberton, J.M.** (2001). Positive genetic correlation between parasite resistance and body size in a free-living ungulate population. *Evolution* **55**: 2116-2125.
- Connan, R.M.** (1969). Studies on the inhibition of development of *Ostertagia* spp. in lambs. *Journal of Helminthology* **43**: 287-292.
- Coop, R.L. and Kyriazakis, I.K.** (2001). Influence of host nutrition on the development and consequences of nematode parasitism in ruminants. *Trends in Parasitology* **17**: 325-330.
- Corn, J.L., Pence, D.B. and Warren, R.J.** (1985). Factors affecting the helminth community structure of adult collared peccaries in southern Texas. *Journal of Wildlife Diseases* **21**: 254-263.
- Cortes, E.** (2002). Incorporating uncertainty into demographic modelling: application to shark populations and their conservation. *Conservation Biology* **16**: 1048-1062.
- Coulson, T., Milner-Gulland, E.J. and Clutton-Brock, T.** (2000). The relative roles of density and climatic variation on population dynamics and fecundity rates in three contrasting ungulate species. *Proceedings of the Royal Society of London Series B* **267**: 1771-1779.
- Courtney, C.H., Parker, C.F., McClure, K.E. and Herd, R.P.** (1984). A comparison of the periparturient rise in faecal egg counts of exotic and domestic ewes. *International Journal for Parasitology* **14**: 377-381.
- Couvillion, C.E.** (1993). Estimation of the numbers of trichostrongylid larvae on pastures. *Veterinary Parasitology* **46**: 197-203.
- Cox, F.E.G.** (2001). Concomitant infections, parasites and immune responses. *Parasitology* **122**: S23-S38.
- Cox, D.D. and Todd, A.C.** (1962). Survey of gastrointestinal parasitism in Wisconsin dairy cattle. *Journal of the American Veterinary Medical Association* **141**: 706-709.
- Cox, J.L., Heyse, J.F. and Tukey, J.W.** (2000). Efficiency estimates of count data that include zero counts. *Experimental Parasitology* **96**: 1-8.
- Coyne, M.J. and Smith, G.** (1992). The development and mortality of the free-living stages of *Haemonchus contortus* in laboratory culture. *International Journal for Parasitology* **22**: 641-650.

**Coyne, M.J. and Smith, G.** (1994). Trichostrongylid parasites of domestic ruminants. In: Scott, M.E. and Smith, G. (Eds.). *Parasitic and Infectious Diseases: epidemiology and ecology*. Academic Press, San Diego. Pp. 235-247.

**Coyne, M.J., Smith, G. and Johnstone, C.** (1991a). Fecundity of gastrointestinal trichostrongylid nematodes of sheep in the field. *American Journal of Veterinary Research* **52**: 1182-1188.

**Coyne, M.J., Smith, G. and Johnstone, C.** (1991b). A study of the mortality and fecundity of *Haemonchus contortus* in sheep following experimental infections. *International Journal for Parasitology* **21**: 847-853.

**Crofton, H.D.** (1948). The ecology of immature phases of trichostrongyle nematodes. I. The vertical distribution of infective larvae of *Trichostrongylus retortaeformis* in relation to their habitat. *Parasitology* **39**: 17-25.

**Crofton, H.D.** (1965). Ecology and biological plasticity of sheep nematodes. I. The effect of temperature on the hatching of eggs of some nematode parasites of sheep. *Cornell Veterinarian* **55**: 242-250.

**Crofton, H.D.** (1971). A model of host-parasite relationships. *Parasitology* **63**: 343-364.

**Crofton, H.D. and Whitlock, J.H.** (1965). Ecology and biological plasticity of sheep nematodes. III. Studies on *Ostertagia circumcincta* (Stadelmann, 1894). *Cornell Veterinarian* **55**: 259-262.

**Crofton, H.D., Whitlock, J.H. and Glazer, R.A.** (1965). Ecology and biological plasticity of sheep nematodes. II. Genetic x environmental plasticity in *Haemonchus contortus* (Rud. 1803). *Cornell Veterinarian* **55**: 251-258.

**Crystal Ball** (2000). *Crystal Ball® 2000 User's Manual*. Decisoneering Inc., Denver, Colorado.

**Dangjin, L.** (1996). Study on a new egg counting technique for *Macracanthorhynchus hirundinaceus* and *Ascaris suum*. *Veterinary Parasitology* **61**: 113-117.

**Das, K.M. and Whitlock, J.H.** (1960). Subspeciation in *Haemonchus contortus* (Rudolphi, 1803) Nematoda, Trichostrongyloidea. *Cornell Veterinarian* **50**: 182-197.

**Daskalov, P.B.** (1972a). *Haemonchus contortus*: attempts to determine biologic differences between the various morphologic types of females. *Experimental Parasitology* **32**: 359-363.

**Daskalov, P.B.** (1972b). *Haemonchus contortus*: factors determining the polymorphism of linguiform females. *Experimental Parasitology* **32**: 364-368.

**Davies, J.E.** (1981). The distribution of *Glossina morsitans* in Botswana from historic to recent times. 17<sup>th</sup> meeting, *International Scientific Council for Trypanosomiasis Research and Control, Arusha, Tanzania, 1981*. Pp. 398-409. (Abstract.)

**Davis, J.W. and Anderson, R.C.** (Eds.) (1971). *Parasitic Diseases of Wild Mammals*. Iowa State University Press, Iowa, USA.

**De Vlas, S.J., Gryseels, B., Van Oortmarsen, G.J., Polderman, A.M. and Habbema, J.D.F.** (1992). A model for variations in single and repeated egg counts in *Schistosoma mansoni* infections. *Parasitology* **104**: 451-460.



- Denisova, S.A.** (Ed.)(1976). *Recommendations for the control of helminths of sheep in Kazakhstan*. Moscow. [In Russian.]
- Denney, R.M.** (1972). Relationships of wildlife to livestock on some developed ranches on the Laikipia Plateau, Kenya. *Range Management* **6**: 415-425. (Abstract.)
- Dezfuli, B.S., Volponi, S., Beltrami, I. And Poulin, R.** (2002). Intra- and interspecific density-dependent effects on growth in helminth parasites of the cormorant, *Phalacrocorax carbo sinensis*. *Parasitology* **124**: 537-544.
- Dias, J.A.T.S., Rosinha, A.J., Santos-Dias, J.A.T. and Travassos-Santos-Dias, J.A.** (1973). Is indiscriminate slaughter of game as a tsetse-control measure justified? (An analysis of the case of Mozambique). *Anais dos Servicos de Veterinaria* 1969-71, **17-19**: 23-53. (Abstract.)
- Diez-Baños, N., Cabaret, J. and Diez-Baños, P.** (1992). Interspecific interactions in naturally acquired nematode communities from sheep abomasum in relation to age of host and season in four areas of Leon (Spain). *International Journal for Parasitology* **22**: 327-334.
- Dineen, J.K., Donald, A.D., Wagland, B.M. and Offner, J.** (1965). The dynamics of the host-parasite relationship. III. The response of sheep to primary infection with *Haemonchus contortus*. *Parasitology* **55**: 515-525.
- Divina, B.P. and Hoglund, J.** (2002), Heterologous transmission with *Dictyocaulus capreolus* from roe deer (*Capreolus capreolus*) to cattle (*Bos taurus*). *Journal of Helminthology* **76**: 125-130.
- Divina, B.P., Wilhelmsson, E., Mattsson, J.G., Waller, P. and Hoglund, J.** (2000). Identification of *Dictyocaulus* spp. in ruminants by morphological and molecular analyses. *Parasitology* **121**: 193-201.
- Dobson, A.P.** (1985). The population dynamics of competition between parasites. *Parasitology* **91**: 317-347.
- Dobson, A.P. and Hudson, P.J.** (1992). Regulation and stability of a free-living host parasite system – *Trichostrongylus tenuis* in Red Grouse. II: Population models. *Journal of Animal Ecology* **61**: 487-498.
- Dobson, A.P. and Hudson, P.J.** (1994). The population biology of *Trichostrongylus tenuis* in red grouse, *Lagopus lagopus scoticus*. Pp. 309-317 in: M.E.Scott and G. Smith (Eds.), *Ecological Epidemiology of Infectious and Parasitic Disease*. Academic Press, New York.
- Dobson, R.J. and Barnes, E.H.** (1995). Interaction between *Ostertagia circumcincta* and *Haemonchus contortus* infection in young lambs. *International Journal for Parasitology* **25**: 495-501.
- Dobson, R.J., Waller, P.J. and Donald, A.D.** (1990). Population dynamics of *Trichostrongylus colubriformis* in sheep: the effect of infection rate on the establishment of infective larvae and parasite fecundity. *International Journal for Parasitology* **20**: 347-352.
- Donald, A.D., Morley, F.H.W., Waller, P.J., Axelsen, A. and Donnelly, J.R.** (1978). Availability to grazing sheep of gastrointestinal nematode infection arising from summer contamination of pastures. *Australian Journal of Agricultural Research* **29**: 189-204.
- Dreyer, K., Fourie, L.J. and Kok, D.J.** (1999). Gastro-intestinal parasites of cattle in the communal grazing system of Botshabelo in the Free State. *Onderstepoort Journal of Veterinary Research* **66**: 145-149.

- Drózdź, J.** (1995). Polymorphism in the Ostertagiinae Lopez-Neyra, 1947 and comments on the systematics of these nematodes. *Systematic Parasitology* **32**: 91-99.
- Dunn, A.** (1968). The wild ruminant as reservoir host of helminth infection. *Symposia of the Zoological Society of London* **24**: 221-248.
- Dunn, A.** (1978). *Veterinary Helminthology*. Frome, Butler and Tanner.
- Dunn, A. and Keymer, A.** (1986). Factors affecting the reliability of the McMaster technique. *Journal of Helminthology* **60**, 260-262.
- Durette-Desset, M.C.** (1982). On generic divisions of the Ostertagiinae (Nematoda). *Annales de Parasitologie Humaine et Comparée* **57**: 375-381. [In French.]
- Durette-Desset, M.C.** (1985). Trichostrongyloid nematodes and their vertebrate hosts: reconstruction of the phylogeny of a parasitic group. *Advances in Parasitology* **24**: 239-306.
- Durette-Desset, M.C.** (1989). Proposed nomenclature for described species in the sub-family Ostertagiinae Lopez-Neyra, 1947. *Annales de Parasitologie Humaine et Comparée* **64**: 356-373. [In French.]
- Durette-Desset, M.C. and Cabaret, J.** (1994). The distances between cuticular ridges follow a Gaussian function in ostertagiine nematodes: the potential use of this phenomenon as a taxonomic criterion. *Systematic Parasitology* **29**: 13-22.
- Durette-Desset, M.C., Hugot, J.P., Darlu, P. and Chabaud, A.G.** (1999). A cladistic analysis of the Trichostrongyloidea (Nematoda). *International Journal for Parasitology* **29**: 1065-1086.
- Eason, G., Coles, C.W. and Gettinby, G.** (1980). *Mathematics and Statistics for the Biosciences*. Ellis Horwood, Chichester.
- Eckert, J. and Hertzberg, H.** (1994). Parasite control in transhumant situations. *Veterinary Parasitology* **54**: 103-125.
- Efron, B. and Tibshirani, R.** (1993). *An Introduction to the Bootstrap*. Chapman and Hall, New York.
- Ehrlich, P.R., Breedlove, D.E., Brussard, P.F. and Sharp, M.A.** (1972). Weather and the regulation of subalpine populations. *Ecology* **53**: 243-247.
- El-Azazy, O.M.E.** (1995). Seasonal changes and inhibited development of the abomasal nematodes of sheep and goats in Saudi Arabia. *Veterinary Parasitology* **58**: 91-98.
- Ellis, T.M., Gregory, A., Turnor, R., Kalkhoven, R.H. and Wroth, R.H.** (1993). Detection of *Haemonchus contortus* surface antigen in faeces from infected sheep. *Veterinary Parasitology* **51**: 85-97.
- Ellis, R.D., Pung, O.J. and Richardson, D.J.** (1999). Site selection by intestinal helminths of the Virginia Opossum (*Didelphis virginiana*). *The Journal of Parasitology* **85**: 1-5.
- Elston, D.A., Moss, R., Boulinier, T., Arrowsmith, C. and Lambin, X.** (2001). Analysis of aggregation, a worked example: numbers of ticks on red grouse. *Parasitology* **122**: 563-569.
- Esch, G.W., Bush, A.O. and Aho, J.M.** (Eds.) (1990). *Parasite Communities: Patterns and Processes*. Chapman and Hall, London.

- Eslami, A., Meydani, M., Maleki, Sh. And Zargarzadeh, A.** (1979). Gastrointestinal nematodes of wild sheep (*Ovis orientalis*) from Iran. *Journal of Wildlife Diseases* **15**: 263-265.
- Eslami, A, Rahbari, S. and Nikbin, S.** (1980). Gastro-intestinal nematodes of gazelle, *Gazella subgutturosa*, in Iran. *Veterinary Parasitology* **7**: 75-78.
- Eysker, M.** (1993). The role of inhibited development in the epidemiology of *Ostertagia* infections. *Veterinary Parasitology* **46**: 259-269.
- Eysker, M. and Kooyman, F.N.J.** (1993). Notes on necropsy and herbage processing techniques for gastrointestinal nematodes of ruminants. *Veterinary Parasitology* **46**: 205-213.
- Eysker, M.** (1997). Some aspects of inhibited development of trichostrongylids in ruminants. *Veterinary Parasitology* **72**: 265-283.
- Fabiyi, J.P. and Copeman, D.B.** (1986). The availability of strongylid larvae to grazing cattle in the wet tropical region of northern Queensland. *Australian Veterinary Journal* **63**: 266-268.
- Fabiyi, J.P., Copeman, D.B. and Hutchinson, G.W.** (1988). Abundance and survival of infective larvae of the cattle nematodes *Cooperia punctata*, *Haemonchus placei* and *Oesophagostomum radiatum* from faecal pats in a wet tropical climate. *Australian Veterinary Journal* **65**: 229-231.
- Fadeev, A.A. and Sludskii, A.A.** (1982). *Saiga in Kazakhstan*. Alma-Ata, Kazakhstan. [In Russian]
- Fakae, B.B. and Chiejina, S.N.** (1988). Further studies on the development and availability of infective larvae of bovine gastrointestinal trichostrongylids on pasture in Eastern Nigeria. *Veterinary Parasitology* **28**: 143-152.
- Familton, A.S.** (1991). Re-examination of gastrointestinal parasite control – the contribution of the ewe. In: *Practical Aspects of Parasite Control. Proceedings of the 21<sup>st</sup> Sheep and Beef Cattle Seminar, Lincoln University, New Zealand, July 1991*. (Ed. Petersen, G.V.), pp. 25-35.
- Farr, M.M. and Luttermoser, G.W.** (1941). Comparative efficiency of zinc sulfate and sugar solutions for the simultaneous flotation of coccidial oocysts and helminth eggs. *The Journal of Parasitology* **27**: 417-424.
- Fine, A.E., Hartman, R., Krecek, R.C. and Groeneveld, H.T.** (1993). Effects of time, from collection to processing, on the recovery of *Haemonchus contortus* third stage larvae from herbage. *Veterinary Parasitology* **51**: 77-83.
- Folstad, I., Nilssen, A.C., Halvorsen, O. and Andersen, O.** (1991). Parasite avoidance – the cause of post-calving migrations in reindeer? *Canadian Journal of Zoology* **69**: 2423-2429.
- Folstad, I., Arneberg, P. and Karter, A.J.** (1996). Antlers and parasites. *Oecologia* **105**: 556-558.
- Forbes, A.B., Huckle, C.A., Gibb, M.J., Rook, A.J. and Nuthall, R.** (2000). Evaluation of the effects of nematode parasitism on grazing behaviour, herbage intake and growth in young grazing cattle. *Veterinary Parasitology* **90**: 111-118.
- Forbes, A.B., Cutler, K.L. and Rice, B.J.** (2002). Sub-clinical parasitism in spring-born, beef suckler calves: epidemiology and impact on growth performance during the first grazing season. *Veterinary Parasitology* **104**: 339-344.

- Foreyt, W.J., Hunter, D., Cook, J.G. and Smith, L.L.** (2000). Susceptibility of elk to lungworms from cattle. *Journal of Wildlife Diseases* **36**: 729-733.
- Fowler, C.W.** (1988). Population dynamics as related to rate of increase per generation. *Evolutionary Ecology* **2**: 197-204.
- Fowler, J., Cohen, L. and Jarvis, P.** (1998). *Practical Statistics for Field Biology* (2<sup>nd</sup> Ed.). Wiley, Chichester.
- Fox, M.T.** (1997). Pathophysiology of infection with gastrointestinal nematodes in domestic ruminants: recent developments. *Veterinary Parasitology* **72**: 285-308.
- France, J., Dhanoa, M.S., Siddons, R.C., Thornley, J.H.M. and Poppi, D.P.** (1988). Estimating the production of faeces by ruminants from faecal marker concentration curves. *Journal of Theoretical Biology* **135**: 383-391.
- Fulford, A.J.C.** (1994). Dispersion and bias: can we trust geometric means? *Parasitology Today* **10**: 446-448.
- Gasbarre, L.C.** (1997). Effects of gastrointestinal nematode infection on the ruminant immune system. *Veterinary Parasitology* **72**: 327-343.
- Gasbarre, L.C., Leighton, E.A. and Davies, C.J.** (1990). Genetic control of immunity to gastrointestinal nematodes of cattle. *Veterinary Parasitology* **37**: 257-272.
- Gasbarre, L.C., Leighton, E.A. and Bryant, D.** (1996). Reliability of a single faecal egg per gram determination as a measure of individual and herd values for trichostrongyle nematodes of cattle. *American Journal of Veterinary Research* **57**: 168-171.
- Gasnier, N., Cabaret, J. and Suarez, V.** (1993). Species and morphs in the Ostertagiinae: An allozyme study of seven species. *International Journal for Parasitology* **23**: 765-770.
- Gatongi, P.M., Gathuma, J.M. and Munyua, W.K.** (1988). Influence of weather on population dynamics of infective larvae of cattle nematodes in Nyeri, Kenya. *Bulletin of Animal Health and Production in Africa* **36**: 148-154.
- Gettinby, G., Bairden, K., Armour, J. and Benitez-Usher, C.** (1979). A prediction model for bovine ostertagiasis. *The Veterinary Record* **105**: 57-59.
- Gettinby, G., Therodoridis, Y. and Whitelaw, A.** (1985). Comparison of two techniques used for the recovery of nematode infective larvae from pasture. *Research in Veterinary Science* **39**: 99-102.
- Giangaspero, M., Bahhady, F.A., Orita, G. and Gruner, L.** (1992). Summer-arrested development of abomasal trichostrongylids in Awassi sheep in semi-arid areas of North-West Syria. *Parasitology Research* **78**: 594-597.
- Gibbons, L.M.** (1979). Revision of the genus *Haemonchus* Cobb, 1898 (Nematoda: Trichostrongylidae). *Systematic Parasitology* **1**: 3-24.
- Gibbons, L.M. and Khalil, L.F.** (1982). A key for the identification of genera of the nematode family Trichostrongylidae Leiper, 1912. *Journal of Helminthology* **56**: 185-233.

- Gibbs, H.C.** (1979). Relative importance of winter survival of larval nematodes in pasture and infected carrier calves in a study of parasitic gastroenteritis in calves. *American Journal of Veterinary Research* **40**: 227-231.
- Gibbs, H.C.** (1986). Hypobiosis in parasitic nematodes: an update. *Advances in Parasitology* **25**: 129-174.
- Gibson, T.E.** (1959). The development and survival of the preparasitic stages of *Nematodirus* spp. on pasture herbage. *Journal of Comparative Medicine and Therapeutics* **68**: 338-344.
- Gibson, T.E.** (1963). Experiments on the epidemiology of nematodiriasis. *Research in Veterinary Science* **4**: 258-268.
- Gibson, T.E. and Everett, G.** (1967). The ecology of the free-living stages of *Trichostrongylus colubriformis*. *Parasitology* **57**: 533-537.
- Gibson, T.E. and Everett, G.** (1972). The ecology of the free-living stages of *Ostertagia circumcincta*. *Parasitology* **64**: 451-460.
- Gibson, T.E. and Everett, G.** (1976). The ecology of the free-living stages of *Haemonchus contortus*. *British Veterinary Journal* **132**: 50-59.
- Gibson, T.E. and Everett, G.** (1981). Ecology of the free-living stages of *Nematodirus battus*. *Research in Veterinary Science* **31**: 323-327.
- Gibson, T.E. and Parfitt, J.W.** (1973). The development of resistance to *Trichostrongylus colubriformis* by lambs under conditions of continuous infection. *Research in Veterinary Science* **15**: 220-223.
- Gibson, T.E. and Whitehead, J.D.** (1981). Changes in the worm burden of lambs under continuous infection with *Ostertagia circumcincta*. *British Veterinary Journal* **137**: 192-195.
- Gill, J.L.** (1990). Uses and abuses of statistical methods in research in parasitology. *Veterinary Parasitology* **36**: 189-209.
- Goldberg, A.** (1968). Development and survival on pasture of gastro-intestinal nematode parasites of cattle. *Journal of Parasitology* **54**: 856-862.
- Goltsman, M., Kruchenkova, E.P. and MacDonald, D.W.** (1996). The Mednyi arcti foxes: treating a population imperilled by disease. *Oryx* **30**: 251-258.
- Gordon, H.M.** (1948). The epidemiology of parasitic diseases, with special reference to studies with nematode parasites of sheep. *Australian Veterinary Journal* **24**: 17-45.
- Gordon, G., O'Callaghan, M. and Tallis, G.M.** (1970). A deterministic model for the life cycle of a class of internal parasites of sheep. *Mathematics in the Biosciences* **8**: 209-226.
- Goskomstat** [State Statistical Committee] (1988). The Economy of Kazakhstan (statistics yearbook). Goskomstat, Almaty. [In Russian]
- Grafen, A. and Woolhouse, M.E.J.** (1993). Does the negative binomial distribution add up? *Parasitology Today* **9**: 475-477.
- Grenfell, B.T.** (1988). Gastrointestinal nematode parasites and the stability and productivity of intensive ruminant grazing systems. *Philosophical Transactions of the Royal Society of London, Series B* **321**: 541-563.

**Grenfell, B.T.** (1992). Parasitism and the dynamics of ungulate grazing systems. *American Naturalist* **139**: 907-929.

**Grenfell, B.T. and Dobson, A.P.** (Eds.) (1995). *Ecology of Infectious Diseases in Natural Populations*. Cambridge University Press.

**Grenfell, B.T. and Gulland, F.M.** (1995). Introduction: ecological impact of parasites on wildlife host populations. *Parasitology* **111**: S3-S14.

**Grenfell, B.T., Smith, G. and Anderson, R.M.** (1986). Maximum-likelihood estimates of the mortality and migration rates of the infective larvae of *Ostertagia ostertagi* and *Cooperia oncophora*. *Parasitology* **92**: 643-652.

**Grenfell, B.T., Smith, G. and Anderson, R.M.** (1987a). The regulation of *Ostertagia ostertagi* populations in calves: the effect of past and current experience of infection on proportional establishment and parasite survival. *Parasitology* **95**: 363-372.

**Grenfell, B.T., Smith, G. and Anderson, R.M.** (1987b). A mathematical model of the population biology of *Ostertagia ostertagi* in calves and yearlings. *Parasitology* **95**: 389-406.

**Grenfell, B.T., Wilson, K., Isham, V.S., Boyd, H.E.G. and Dietz, K.** (1995). Modelling patterns of parasite aggregation in natural populations: trichostrongylid nematode-ruminant interactions as a case study. *Parasitology* **111**: S135-S151.

**Grenfell, B.T., Amos, W., Arneberg, P., Bjørnstad, O.N., Greenman, J.V., Harwood, J., Lanfranchi, P., McLean, A.R., Norman, R.A., Read, A.F. and Skorping, A.** (2002). Visions for future research in wildlife epidemiology. In: Hudson, P.J., Rizzoli, A., Grenfell, B.T., Heesterbeek, H. and Dobson, A. (Eds.). *The Ecology of Wildlife Diseases*. Oxford University Press. Pp. 151-164.

**Gruner, L., Berbigier, P., Cortet, J. and Sauve, C.** (1989). Effects of irrigation on appearance and survival of infective larvae of goat gastro-intestinal nematodes in Guadeloupe (French West Indies). *International Journal for Parasitology* **19**: 409-415.

**Gruzdev, A.R. and Pronyaev, A.V.** (1994). Determination of the age of saiga antelopes from cement strata of teeth. *Zoologicheskii Zhurnal* **73**: 223-226. [In Russian.]

**Gulland, F.M.D.** (1992). The role of nematode parasites in Soay sheep (*Ovis aries* L.) mortality during a population crash. *Parasitology* **105**: 493-503.

**Gulland, F.M.D. and Fox, M.** (1992). Epidemiology of nematode infections of Soay sheep (*Ovis aries* L.) on St. Kilda. *Parasitology* **105**: 481-492.

**Gulland, F.M.D., Albon, S.D., Pemberton, J.M., Moorcroft, P.R. and Clutton-Brock, T.H.** (1993). Parasite-associated polymorphism in a cyclic ungulate population. *Proceedings of the Royal Society of London Series B* (1993): 7-13.

**Guyatt, H.L., Brooker, S. and Donnelly, C.A.** (1999). Can prevalence of infection in school-aged children be used as an index of community prevalence? *Parasitology* **118**: 257-268.

**Hall, A.** (1982). Intestinal helminths of man: the interpretation of egg counts. *Parasitology* **85**: 605-613.

- Halvorsen, O.** (1986). On the relationship between social status of host and risk of parasitic infection. *Oikos* **47**: 71-74.
- Halvorsen, O., Stien, A., Irvine, J., Langvatn, R. and Albon, S.** (1999). Evidence for continued transmission of parasitic nematodes in reindeer during the Arctic winter. *International Journal for Parasitology* **29**: 567-579.
- Hanski, I.** (1991). Single species metpopulation dynamics: Concepts, models and observations. *Biological Journal of the Linnean Society* **42**: 17-38.
- Harvell, C.D., Mitchell, C.E., Ward, J.R., Alitzer, S., Dobson, A.P., Ostfeld, R.S. and Samuel, M.D.** (2002). Climate warming and disease risks for terrestrial and marine biota. *Science* **296**: 2158-2162.
- Hassell, M.P., Lawton, J.H. and May, R.M.** (1976). Patterns of dynamical behaviour in single-species populations. *Journal of Animal Ecology* **45**: 471-486.
- Hassell, M.P., Comins, H.N. and May, R.M.** (1991). Spatial structure and chaos in insect population dynamics. *Nature* **353**: 255-258.
- Haukisalmi, V., Henttonen, H. and Vikman, P.** (1996). Variability of sex ratio, mating probability and egg production in an intestinal nematode in its fluctuating host population. *International Journal for Parasitology* **26**: 755-764.
- Heinichen, I.G.** (1973). Parasitological studies on Impala: preliminary report. *Journal of the South African Veterinary Association* **44**: 265-269.
- Helle, O.** (1971). The survival of nematodes and cestodes of sheep in the pasture during the winter in eastern Norway. *Acta Veterinaria Scandinavica* **12**: 504-512.
- Henriksen, S.A., Bentholt, B.R. and Nielsen-Englyst, A.** (1976). Investigations concerning bovine gastro-intestinal strongyles. II. Seasonal variations in the herbage infestation with infective larvae. *Nordskaer Veterinarmedicin* **28**: 201-209. [In Norwegian]
- Herlich, H.** (1954). The life history of *Nematodirus helvetianus* May, 1920, a nematode parasitic in cattle. *Journal of Parasitology* **40**: 60-70.
- Herlich, H., Gasbarre, L.C. and Douvres, F.W.** (1984). Infectivity and pathogenicity of three isolates of *Ostertagia ostertagi* in cattle. *Veterinary Parasitology* **16**: 253-260.
- Hess, G.** (1996). Disease in metapopulation models: implications for conservation. *Ecology* **77**: 1617-1632.
- Hilborn, R. and Mangel, M.** (1997). *The Ecological Detective: Confronting Models with Data*. Monographs in Population Biology 28. Princeton University Press, New Jersey.
- Hoberg, E.P. and Lichtenfels, J.R.** (1994). Phylogenetic systematic analysis of the Trichostrongylidae (Nematoda), with an initial assessment of coevolution and biogeography. *The Journal of Parasitology* **80**: 976-996.
- Hoberg, E.P., Monsen, K.J., Kutz, S. and Blouin, M.S.** (1999). Structure, biodiversity, and historical biogeography of nematode faunas in holarctic ruminants: morphological and molecular diagnoses for *Teladorsagia boreoarticus* n. sp. (Nematoda: Ostertagiinae), a dimorphic cryptic species in muskoxen (*Ovibos moschatus*). *Journal for Parasitology* **85**: 910-934.

- Hoberg, E.P., Kocan, A.A. and Rickard, L.G.** (2001). Gastrointestinal strongyles in wild ruminants. In: *Parasitic Diseases of Wild Mammals* (Eds. Samuel, W.M., Pybus, M.J. and Kocan, A.A.). Iowa State University Press, Ames. Pp. 193-227.
- Holland, C.V. and Cox, D.M.** (2001). *Toxocara* in the mouse: a model for parasite-altered host behaviour? *Journal of Helminthology* **75**: 125-135.
- Holmes, J.C.** (1961). Effects of concurrent infections on *Hymenolepis diminuta* (Cestoda) and *Moniliformis dubius* (Acanthocephala). I. General effects and comparison with crowding. *The Journal of Parasitology* **47**: 209-216.
- Holmes, J.C.** (1962). Effects of concurrent infections on *Hymenolepis diminuta* (Cestoda) and *Moniliformis dubius* (Acanthocephala). III. Effects in hamsters. *The Journal of Parasitology* **48**: 97-100.
- Hoodless, A.N., Kurtenbach, K., Nuttall, P.A. and Randolph, S.E.** (2002). The impact of ticks on pheasant territoriality. *Oikos* **96**: 245-250.
- Horak, I.G.** (1978). Parasites of domestic and wild animals in South Africa. X. Helminths in Impala. *Onderstepoort Journal of Veterinary Research* **45**: 221-228.
- Horak, I.G.** (1981). Host specificity and the distribution of the helminths of sheep, cattle, impala and blesbok according to climate. *Journal of the South African Veterinary Association* **52**: 201-206.
- Horak, I.G., De Vos, V. and Brown, M.** (1983). Parasites of domestic and wild animals in South Africa. XVI. Helminth and arthropod parasites of blue and black wildebeest (*Connochaetes taurinus* and *Connochaetes gnou*). *Onderstepoort Journal of Veterinary Research* **50**: 243-255.
- Horak, I.G., MacIvor, K.M.de F. and Greeff, C.J.** (2001). Parasites of domestic and wild animals in South Africa. XXXIX. Helminth and arthropod parasites of Angora goats in the southern Karoo. *Onderstepoort Journal of Veterinary Research* **68**: 27-35.
- Hoste, H.** (2001). Adaptive physiological processes in the host during gastrointestinal parasitism. *International Journal for Parasitology* **31**: 231-244.
- Hoste, H., Leveque, H. and Dorchies, Ph.** (2001). Comparison of nematode infections of the gastrointestinal tract in Angora and dairy goats in a rangeland environment: relations with the feeding behaviour. *Veterinary Parasitology* **101**: 127-135.
- Houdijk, J.G.M., Kyriazakis, I., Coop, R.L. and Jackson, F.** (2001). The expression of immunity to *Teladorsagia circumcincta* in ewes and its relationship to protein nutrition depend on body protein reserves. *Parasitology* **122**: 661-672.
- Howard, S.C., Donnelly, C.A. and Chan, M.-S.** (2001). Methods for estimation of associations between multiple species parasite infections. *Parasitology* **122**: 233-251.
- Hsu, C.K. and Levine, N.D.** (1977). Degree-day concept in development of infective larvae of *Haemonchus contortus* and *Trichostrongylus colubriformis* under constant and cyclic conditions. *American Journal of Veterinary Research* **138**: 1115-1119.
- Hudson, P.J. and Dobson, A.P.** (1995). Macroparasites: observed patterns. In: Grenfell, B.T. and Dobson, A.P. (1995). *Ecology of Infectious Diseases in Natural Populations*. Cambridge University Press.



- Hudson, P.J., Dobson, A.P. and Newborn, D.** (1998). Prevention of parasite cycles by parasite removal. *Science* **282**: 2256-2258.
- Hudson, P.J., Rizzoli, A., Grenfell, B.T., Heesterbeek, H. and Dobson, A.P.,** (Eds.) (2002). *The Ecology of Wildlife Diseases*. Oxford University Press. 197pp.
- Humbert J.F. and Cabaret, J.** (1995). Use of random amplified polymorphic DNA for identification of ruminant trichostrongylid nematodes. *Parasitology Research* **81**:1-5.
- Hunter, G.C. and Quenouille, M.H.** (1952). A statistical examination of the worm egg count sampling technique for sheep. *Journal of Helminthology* **4**: 157-170.
- Hunter, C.M., Moller, H. and Fletcher, D.** (2000). Parameter uncertainty and elasticity analyses of a population model: setting research priorities for shearwaters. *Ecological Modelling* **134**: 299-324.
- Hutchings, M.R., Kyriazakis, I., Anderson, D.H., Gordon, I.J. and Coop, R.L.** (1998). Behavioural strategies used by parasitized and non-parasitized sheep to avoid ingestion of gastro-intestinal nematodes associated with faeces. *Animal Science* **67**: 97-106.
- Inglis, W.G.** (1965). Patterns of evolution in parasitic nematodes. Pp. 79-124 in: Taylor, A.E.R. (Ed.). *Evolution of Parasites*. 3<sup>rd</sup> Symposium of the British Society for Parasitology. Blackwell Scientific, Oxford.
- Irgashev, I. Kh.** (1973). *Helminths and Helminthoses of the Karakul sheep*. National Academy of Sciences of the Uzbekistan Soviet Socialist Republic, Institute of Zoology and Parasitology, Tashkent, Uzbekistan. 284pp. [In Russian]
- Irvine, R.J., Stien, A., Halvorsen, O., Langvatn, R. and Albon, S.D.** (2000). Life-history strategies and population dynamics of abomasal nematodes in Svalbard reindeer (*Rangifer tarandus platyrhynchus*). *Parasitology* **120**: 297-311.
- Irvine, R.J., Stien, A., Dallas, J.F., Halvorsen, O., Langvatn, R. and Albon, S.D.** (2001). Contrasting regulation of fecundity in two abomasal nematodes of Svalbard reindeer (*Rangifer tarandus platyrhynchus*). *Parasitology* **122**: 673-681.
- Isham, V.** (1995). Stochastic models of host-macroparasite interaction. *The Annals of Applied Probability* **5**: 720-740.
- Jackson, F. and Christie, M.G.** (1979). Observations on the egg output from continuous low level infections with *Ostertagia circumcincta* in lambs. *Research in Veterinary Science* **27**: 244-245.
- Jacquiet, Ph., Humbert, J.F., Comes, A.M., Thiam, A. and Chiekh, D.** (1995a). Ecological, morphological and genetic characterization of sympatric *Haemonchus* spp. parasites of domestic ruminants in Mauritania. *Parasitology* **110**: 483-492.
- Jacquiet, Ph., Coloas, F., Cabaret, J., Dia, M.L., Chiekh, D. and Thiam, E.** (1995b). Dry areas: an example of seasonal evolution of helminth infection of sheep and goats in southern Mauritania. *Veterinary Parasitology* **56**: 137-148.
- Jacquiet, Ph., Cabaret, J., Chiekh, D. and Thiam, E.** (1995c). Experimental study of survival strategy of *Haemonchus contortus* in sheep during the dry season in desert areas of the Mauritania. *Journal of Parasitology* **81**: 1013-1015.

- Jacquet, Ph., Cabaret, J., Dia, M.L., Chiekh, D. and Thiam, E.** (1996). Adaptation to arid environment: *Haemonchus longistipes* in dromedaries of Saharo-Sahelian areas of Mauritania. *Veterinary Parasitology* **66**: 193-204.
- Jacquet, Ph., Cabaret, J., Chiekh, D. and Thiam, E.** (1997). Identification of *Haemonchus* species in domestic ruminants based on morphometrics of spicules. *Parasitology Research* **83**: 82-86.
- Jacquet, P., Cabaret, J., Thiam, A. and Cheikh, D.** (1998). Host range and the maintenance of *Haemonchus* spp. in an adverse arid climate. *International Journal for Parasitology* **28**: 253-261.
- Jauniaux, T. and Coignoul, F.** (2001). Morbillivirus in common seals stranded off the coasts of Belgium and northern France during summer 1998. *The Veterinary Record* **148**: 587-591.
- Jiang-Zhigang, Feng-Zuojian, Wang-Zuwang, Chen-Liwei, Cai-Ping and Li-Yongbo** (1995). Historical and current distributions of Przewalski's gazelles. *Acta Theriologica Sinica* **15**: 241-245. (Abstract.)
- Kabasa, J.D., Opuda-Asibo, J. and ter Meulen, U.** (2000). The effect of oral administration of polyethylene glycol on faecal helminth egg counts in pregnant goats grazed on browse containing condensed tannins. *Tropical Animal Health and Production* **32**: 73-86.
- Kao, R.R., Leathwick, D.M., Roberts, M.G. and Sutherland, I.A.** (2000). Nematode parasites of sheep: a survey of epidemiological parameters and their application in a simple model. *Parasitology* **121**: 85-103.
- Karabaev, D.K.** (1953). Changes in the helminth fauna of sheep transferred to Betpak-Dala (central Kazakhstan). Pp. 284-287 in: *Works in Helminthology in Honour of the 75<sup>th</sup> Anniversary of Academician K.I. Skrjabin*. Academy of Sciences of the USSR, Moscow. [In Russian.]
- Karabaev, D.K.** (1973). Zonal distribution of helminths and helminthoses of sheep in Kazakhstan. Pp. 283-294 in: *Chemoprophylaxis, pathogenesis and epizootology of helminthoses of domestic animals. Transactions of the Kazakhstan Scientific Research Veterinary Institute. Vol. 15*. Almaty, Kazakhstan. [In Russian.]
- Keeling, M.J.** (1999). Spatial models of interacting populations. In: *Advanced Ecological Theory. Principles and Applications*. (Ed. McGlade, J.). Blackwell Science, Oxford. Pp. 64-99.
- Keeling, M.J.** (2000). Metapopulation moments: coupling, stochasticity and persistence. *Journal of Animal Ecology* **69**: 725-736.
- Keeling, M.J., Wilson, H.B. and Pacala, S.W.** (2002). Reinterpreting space, time lags, and functional responses in ecological models. *Science* **290**: 1758-1761.
- Keith, L.B., Cary, J.R., Yuill, T.M. and Keith, I.M.** (1985). Prevalence of helminths in a cyclic snowshoe hare population. *Journal of Wildlife Diseases* **21**: 233-253.
- Kennedy, C.R.** (1975). *Ecological Animal Parasitology*. Blackwell Scientific Publications, Oxford.
- Kennedy, S.** (1990). A review of the 1988 European seal morbillivirus epidemic. *The Veterinary Record* **127**: 563-567.

- Keymer, A.** (1982). Density-dependent mechanisms in the regulation of intestinal helminth populations. *Parasitology* **84**: 573-587.
- Kingsbury, P.A.** (1965). Relationship between egg counts and worm burdens of young sheep. *The Veterinary Record* **77**: 900-901.
- Kinnear, P.R. and Gray, C.D.** (2000). *SPSS for Windows Made Simple. Release 10*. Psychology Press, Hove.
- Kistner, T.P., Matlock, S.M., Wyse, D. and Mason, G.E.** (1977). Helminth parasites of bighorn sheep in Oregon. *Journal of Wildlife Diseases* **13**: 125-130.
- Krecek, R.C., Murrell, K.D. and Douglass, L.W.** (1990). Effects of microclimatic variables on the availability and movement of third-stage larvae of *Ostertagia ostertagi* on herbage. *Onderstepoort Journal of Veterinary Research* **57**: 133-135.
- Krecek, R.C., Groeneveld, H.T. and Maritz, J.I.** (1992). A preliminary study of the effect of microclimate on third-stage larvae of *Haemonchus contortus* and *Haemonchus placei* on irrigated pasture. *International Journal for Parasitology* **22**: 747-752.
- Kremer, J.N.** (1983). Ecological implications of parameter uncertainty in stochastic simulation. *Ecological Modelling* **18**: 187-207.
- Kruuk, L.E.B., Clutton-Brock, T.H., Albon, S.D., Pemberton, J.M. and Guinness, F.E.** (1999). Population density affects sex ratio variation in red deer. *Nature* **399**: 459-461.
- Kunz, W.** (2002). When is a parasite species a species? *Trends in Parasitology* **18**: 121-124.
- Kuznetsov, V.I. and Dikov, G.I.** (1979). The situation of helminths of sheep in Kazakhstan and means of prophylaxis against the most pathogenic. Pp. 96-100 and tables in: Zadirozhiy, I.F. (Ed.). *Proceedings of the Republic seminar on the control of parasitic diseases of domestic animals in commemoration of the hundredth anniversary of the birth of academician K.I. Skryabin*. Scientific Veterinary Research Institute, Almaty, Kazakhstan. [In Russian]
- Kyriazakis, I., Tolkamp, B.J. and Hutchings, M.R.** (1998). Towards a functional explanation of the occurrence of anorexia during parasitic infections. *Animal Behaviour* **56**: 265-274.
- Lancaster, M.B.** (1970). The recovery of infective nematode larvae from herbage samples. *Journal of Helminthology* **44**: 219-230.
- Lane, C.** (1928). The mass diagnosis of hookworm infection. *Ibid.* May supplement, 1-148. Cited in Stoll (1930).
- Lavrov, L.I.** (1970). Shared helminths of wild and domestic ruminants in south Kazakhstan. *Contributions to the Natural Nidality of Diseases* **3**: 134-138. [In Russian.]
- Le Jambre, L.F.** (1995). Relationship of blood loss to worm numbers, biomass and egg production in *Haemonchus* infected sheep. *International Journal for Parasitology* **25**: 269-273.
- Le Jambre, L.F. and Ratcliffe, L.H.** (1976). Response of *Haemonchus contortus cayugensis* to a change in the ratio of smooth to linguiform. *Parasitology* **73**: 213-222.
- Le Jambre, L.F. and Whitlock, J.H.** (1976). Changes in the hatch rate of *Haemonchus contortus* eggs between geographic regions. *Parasitology* **73**: 223-238.

- Leathwick, D.M., Miller, C.M., Brown, A.E. and Sutherland, I.A.** (1999). The establishment rate of *Ostertagia circumcincta* and *Trichostrongylus colubriformis* in lactating Romney ewes. *International Journal for Parasitology* **29**: 315-320.
- Levine, N.D.** (1963). Weather, climate, and the bionomics of ruminant nematode larvae. *Advances in Veterinary Science* **8**: 215-261.
- Levine, N.D.** (1980). *Nematode Parasites of Domestic Animals and of Man*. Burgess, Minneapolis.
- Levine, N.D., Krishna, N.M., Clark, D.T. and Aves, I.J.** (1960). A comparison of nematode egg counting techniques for cattle and sheep feces. *American Journal of Veterinary Research*, May 1960: 511-515.
- Levine, N.D., Todd, K.S. and Boatman, P.A.** (1974). Development and survival of *Haemonchus contortus* on pasture. *American Journal of Veterinary Research* **35**: 1413-1422.
- Liang, S., Maszle, D. and Spear, R.C.** (2002). A quantitative framework for a multi-group model of *Schistosomiasis japonicum* transmission dynamics and control in Sichuan, China. *Acta Tropica* **82**: 263-277.
- Lichtenfels, J.R., Hoberg, E.P.** (1993). The systematics of nematodes that cause ostertagiasis in domestic and wild ruminants in North America: an update and a key to species. *Veterinary Parasitology* **46**: 33-53.
- Lichtenfels, J.R., Hoberg, E.P. and Zarlenga, D.S.** (1997). Systematics of gastrointestinal nematodes of domestic ruminants: advances between 1992 and 1995 and proposals for future research. *Veterinary Parasitology* **72**: 225-245.
- Lindenmayer, D.B. and Lacy, R.C.** (2002). Small mammals, habitat patches and PVA models: a field test of model predictive ability. *Biological Conservation* **103(3)**: 247-265.
- Lively, C.M., and Apanius, V.** (1995). Genetic diversity in host-parasite interactions. In: Grenfell, B.T. and Dobson, A.P. (Eds.). *Ecology of Infectious Diseases in Natural Populations*. Cambridge University Press. Pp. 421-449.
- Lloyd, S.** (1995). Environmental influences on host immunity. In: Grenfell, B.T. and Dobson, A.P. (Eds.). *Ecology of Infectious Diseases in Natural Populations*. Publications of the Newton Institute No. 7. Cambridge University Press. Pp. 327-361.
- Lundervold, M.** (2001). Infectious diseases of saiga antelopes and domestic livestock in Kazakhstan. PhD Thesis, University of Warwick.
- Lundervold, M., Langvatn, R. and Milner-Gulland, E.J.** (2003). A comparison of age determination methods for the saiga antelope (*Saiga tatarica*). *Wildlife Biology (In press)*.
- Lushchekina, A.A., Neronov, V.M., Valery, M., Amgalan, L. and Dulamtseren, S.** (1999). The status and prospects for conservation of the Mongolian saiga *Saiga tatarica mongolica*. *Oryx* **33**: 20-29.
- MAFF** (1984). *Manual of Veterinary Investigation Laboratory Techniques*. Reference Book 390. 3<sup>rd</sup> ed. HMSO, London.
- MAFF** (1986). *Manual of Veterinary Parasitological Laboratory Techniques*. Reference Book 418. 3<sup>rd</sup> ed. HMSO, London. 160pp.

- Mahoney** (1968). *Laboratory Techniques in Zoology*. Butterworths, London.
- Manton, V.J.A., Peacock, R., Poynter, D., Silverman, P.H. and Terry, R.J.** (1962). The influence of age on naturally acquired resistance to *Haemonchus contortus* in lambs. *Research in Veterinary Science* **3**: 308-314.
- Martin, P.J., Le Jambre, L.F. and Claxton, J.H.** (1981). The impact of refugia on the development of thiabendazole resistance in *Haemonchus contortus*. *International Journal for Parasitology* **11**: 35-41.
- May, R.M.** (1975). Biological populations obeying difference equations: stable points, limit cycles, and chaos. *Journal of Theoretical Biology* **51**: 511-524.
- May, R.M. and Anderson, R.M.** (1978). Regulation and stability of host-parasite population interactions. II. Destabilizing processes. *Journal of Animal Ecology* **47**: 249-267.
- May, R.M. and Anderson, R.M.** (1979). Population biology of infectious diseases: Part II. *Nature* **280**: 455-461.
- Medica, D.L. and Sukhdeo, M.V.K.** (2001). Estimating transmission potential in gastrointestinal nematodes (Order: Strongylida). *The Journal of Parasitology* **87**: 442-445.
- Medley, G.F.** (2002). The epidemiological consequences of optimisation of the individual host immune response. *Parasitology* **125**: S61-S70.
- Mes, T.H.M., Ploeger, H.W., Terlouw, M., Kooyman, F.N.J., Van der Ploeg, M.P.J. and Eysker, M.** (2001). A novel method for the isolation of gastro-intestinal nematode eggs that allows automated analysis of digital images of egg preparations and high throughput screening. *Parasitology* **123**: 309-314.
- Michel, J.F.** (1968). Faecal egg counts in infections of gastro-intestinal nematodes in cows. *The Veterinary Record*, Feb. 3<sup>rd</sup>, 132-133.
- Michel, J.F.** (1969a). Observations on the faecal egg count of calves naturally infected with *Ostertagia ostertagi*. *Parasitology* **59**: 829-835.
- Michel, J.F.** (1969b). The regulation of egg output by *Ostertagia ostertagi* in calves infected once only. *Parasitology* **59**: 767-774.
- Michel, J.F.** (1974). Arrested development of nematodes and some related phenomena. *Advances in Parasitology* **12**: 279-366.
- Michel, J.F., Lancaster, M.B. and Hong, C.** (1972). Host induced effects on the vulval flap of *Ostertagia ostertagi*. *International Journal for Parasitology* **2**: 305-317.
- Michel, J.F., Lancaster, M.B. and Hong, C.** (1973). *Ostertagia ostertagi*: protective immunity in calves. *Experimental Parasitology* **33**: 179-186.
- Milner-Gulland, E.J.** (1994). A population model for the management of the saiga antelope. *Journal of Applied Ecology* **31**: 25-39.
- Milner-Gulland, E.J., Kholodova, M.V., Bekenov, O.M., Grachev, Iu.A., Amgalan, L. and Lushchekina, A.A.** (2001). Dramatic declines in saiga antelope populations. *Oryx* **35**: 340-345.
- MLC** (1983). *Feeding the ewe*. Meat and Livestock Commission, Milton Keynes. 26pp.

- Montgomery, S.S.J. and Montgomery, W.I.** (1988). Cyclic and non-cyclic dynamics in populations of the helminth parasites of wood mice, *Apodemus sylvaticus*. *Journal of Helminthology* **62**: 78-90.
- Mordi, A.R.** (1989). The future of animal wildlife and its habitat in Botswana. *Environmental Conservation* **16**: 147-156. (Abstract.)
- Moriya, S.** (1954). The reliability of the current diagnostic methods for the identification of helminth eggs. *Parasitology* **44**: 300-303.
- Mukhmetalin, K.D. and Tursunkulov, Sh.Zh.** (1975). On the age and seasonal dynamics of helminthoses of sheep and timing of anthelmintic treatment in south Kazakhstan. Pp. 238-239 in: *Diseases of Sheep and Means of Their Control in Kazakhstan*. Kazakhstan Scientific Research Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]
- Murdoch, W.W. and Briggs, C.J.** (2002). Spatial dynamics of measles epidemics. *Trends in Ecology and Evolution* **17**: 399-401.
- Murdoch, W.W. and Oaten, A.** (1975). Predation and population stability. *Advances in Ecological Research* **9**: 1-131.
- Murray, D.L., Cary, J.R. and Lloyd, B.K.** (1997). Interactive effects of sublethal nematodes and nutritional status on snowshoe hare vulnerability to predation. *Journal of Animal Ecology* **66**: 250-264.
- Murray, D.L., Lloyd, B.K. and Cary, J.R.** (1998). Do parasitism and nutritional status interact to affect production in snowshoe hares? *Ecology* **79**: 1209-1222.
- Murrell, K.D., Leighton, E.A., Boswell, B.A. and Gasbarre, L.C.** (1989). Comparison of egg excretion and serum pepsinogen levels as measures of nematode worm burdens in calves with limited pasture exposure. *The Journal of Parasitology* **75**: 360-366.
- Mustafin, A.O.** (1987). The helminthosis situation on sheep farms in Pavlodar oblast. Pp. 64-69 in: *Problems of Veterinary Parasitology in Kazakhstan. A Collection of Scientific Transactions*. Kazakhstan Scientific Investigation Veterinary Institute, Almaty. [In Russian.]
- McFarlane, R.G.** (1997). Immunity to nematodes in ruminants. In: Barrell (ed.). *Sustainable Control of Internal Parasites in Ruminants*. Lincoln University, New Zealand.
- McKenna, P.B.** (1973). The effect of storage on the infectivity and parasitic development of third stage *Haemonchus contortus* larvae in sheep. *Research in Veterinary Science* **14**: 312-316.
- McKenna, P.B.** (1981). The diagnostic value and interpretation of faecal egg counts in sheep. *New Zealand Veterinary Journal* **29**: 129-132.
- McKenna, P.B.** (1990). The effect of previous cold storage on the subsequent recovery of infective third stage nematode larvae from sheep faeces. *Veterinary Parasitology* **80**: 167-172.
- Nansen, P., Foldager, J., Hansen, J.W., Henriksen, Sv.Aa. and Jørgensen, R.J.** (1988). Grazing pressure and acquisition of *Ostertagia ostertagi* in calves. *Veterinary Parasitology* **27**: 325-335.

**Nekipelova, R.A., Papikhina, K.M., Gordeev, Yu.S. and Aringazin, S.** (1975). The influence of external surroundings on the development of eggs of *Nematodirus* in winter conditions. Pp. 241-242 in: *Diseases of Sheep and Means of Their Control in Kazakhstan*. Kazakhstan Scientific Research Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]

**Nicholls, J. and Obendorf, D.L.** (1994). Application of a composite faecal egg count procedure in diagnostic parasitology. *Veterinary Parasitology* **52**: 337-342.

**Niezen, J.H., Charleston, W.A.G., Hodgson, J., Mackay, A.D. and Leathwick, D.M.** (1996). Controlling internal parasites in grazing ruminants without recourse to anthelmintics: approaches, experiences and prospects. *International Journal for Parasitology* **26**: 983-992.

**Niezen, J.H., Charleston, W.A.G., Hodgson, J., Miller, C.M., Waghorn, T.S. and Robertson, H.A.** (1998a). Effect of plant species on the larvae of gastrointestinal nematodes which parasitise sheep. *International Journal for Parasitology* **28**: 791-803.

**Niezen, J.H., Miller, C.M., Robertson, H.A., Wilson, S.R. and Mackay, A.D.** (1998b). Effect of topographical aspect and farm system on the population dynamics of *Trichostrongylus* larvae on a hill pasture. *Veterinary Parasitology* **78**: 37-48.

**Niezen, J.H., Charleston, W.A.G., Hodgson, J., Miller, C.M., Waghorn, T.S. and Robertson, H.A.** (1998c). Establishment and fecundity of *Ostertagia circumcincta* and *Trichostrongylus colubriformis* in lambs fed lotus (*Lotus pedunculatus*) or perennial ryegrass (*Lolium perenne*). *Veterinary Parasitology* **78**: 13-21.

**Nikitenko, G.I.** (1968). Means of control of helminthoses of domestic animals in Dzhambul oblast. In: *Proceedings of the Advisory Seminar on the Control of Helminthoses of Domestic Animals*. Kazakhstan Scientific Investigation Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. Pp. 95-97. [In Russian.]

**Nilsson, O.** (1971). The inter-relationship of endo-parasites in wild cervids (*Capreolus capreolus* L. and *Alces alces* L.) and domestic ruminants in Sweden. *Acta Veterinaria Scandinavica* **12**: 36-68.

**Northam, J.I. and Rocha, U.F.** (1958). On the statistical analysis of worm counts in chickens. *Experimental Parasitology* **7**: 428-438.

**Noyes, H.A., Ambrose, P., Barker, F., Begon, M., Bennett, M., Bown, K.J. and Kemp, S.J.** (2002). Host specificity of *Trypanosoma* (*Herpetosoma*) species: Evidence that bank voles (*Clethrionomys glareolus*) carry only one *T.(H.) evotomys* 18S rRNA genotype but wood mice (*Apodemus sylvaticus*) carry at least two polyphyletic parasites. *Parasitology* **124**: 185-190.

**Nwosu, C.O., Ogunrinade, A.F. and Fagbemi, B.O.** (1996). The seasonal prevalence of *Haemonchus* species in Red Sokoto (Maradi) goats in Nigeria. *Veterinary Research Communications* **20**: 367-370.

**O'Reagain, P.J. and Schwartz, J.** (1995). Dietary selection and foraging strategies of animals on rangeland. Coping with spatial and temporal variability. In: *Recent developments in the nutrition of herbivores. Proceedings of the IVth International Symposium on the Nutrition of Herbivores, Clermont-Ferrand (France), Sept 1995* (Eds. Journet, M., Grenet, E., Farce, M.H., Thierez, M. and Demarquilly, C.), pp. 407-423.

**Ogonsusi, R.A.** (1979). Pasture infectivity with trichostrongylid larvae in the northern Guinea Savanna of Nigeria. *Research in Veterinary Science* **26**: 320-323.

- Okon E.D. and Enyenihi, U.K.** (1977). Development and survival of *Haemonchus contortus* larvae on pastures in Ibadan. *Tropical Animal Health and Production* **9**: 7-10.
- Omarov, B.B.** (1996a). The influence of the external surroundings on the development of eggs and larvae of *Nematodirus* in semidesert and hilly zones. Pp. 44-49 in: *Invasive Diseases of Domestic Animals and Their Prophylaxis. A Collection of Scientific Transactions*. Kazakhstan Scientific Investigation Veterinary Institute, Kazakhstan Academy of Agricultural Science, Almaty. [In Russian.]
- Omarov, B.B.** (1996b). Nematodirosis of sheep in the south-east and means of control. Pp. 49-56 in: *Invasive Diseases of Domestic Animals and Their Prophylaxis. A Collection of Scientific Transactions*. Kazakhstan Scientific Investigation Veterinary Institute, Kazakhstan Academy of Agricultural Science, Almaty. [In Russian.]
- Onar, E.** (1975). Observations on *Nematodirus abnormalis* (May, 1920): isolation, eggs and larvae, pre-parasitic development. *British Veterinary Journal* **131**: 231-239.
- Onyali, I.O., Onwuliri, C.O.E. and Ajayi, J.A.** (1990). Development and survival of *Haemonchus contortus* larvae on pasture at Vom, Plateau State, Nigeria. *Veterinary Research Communications* **14**: 211-216.
- Orinbaev, B.** (1968). Intestinal cestodes and infectious enterotoxaemia. In: *Proceedings of the Advisory Seminar on the Control of Helminthoses of Domestic Animals*. Kazakhstan Scientific Investigation Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. Pp. 97-98. [In Russian.]
- Osipov, P.P., Karamendin, O.S. and Gubaidulin, N.A.** (1968). Nematodirosis of sheep and methods for its control. Pp. 102-103 in: *Proceedings of The Advisory Seminar on the Control of Helminthoses of Domestic Animals*. Kazakhstan Scientific Investigation Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]
- Pacala, S.W. and Dobson, A.P.** (1988). The relation between the number of parasites/host and host age: population dynamic causes and maximum likelihood estimation. *Parasitology* **96**: 197-210.
- Pandey, V.S.** (1972). Effect of temperature on survival of the free-living stages of *Ostertagia ostertagi*. *Journal of Parasitology* **58**: 1042-1046.
- Pandey, V.S., Chaer, A. and Dakkak, A.** (1989). Effect of temperature on development of the free-living stages of *Ostertagia circumcincta*. *Veterinary Parasitology* **32**: 193-197.
- Pandey, V.S., Chitate, F. and Nyanzunda, T.M.** (1993). Epidemiological observations on gastro-intestinal nematodes in communal land cattle from the highveld of Zimbabwe. *Veterinary Parasitology* **51**: 99-106.
- Papadopoulos, E., Himonas, C. and Coles, G.C.** (2001). Drought and flock isolation may enhance the development of anthelmintic resistance in nematodes. *Veterinary Parasitology* **97**: 253-259.
- Parkin, J.T.** (1976). The effect of moisture supply upon the development and hatching of the eggs of *Nematodirus battus*. *Parasitology* **73**: 343-354.
- Paton, G., Thomas, R.J. and Waller, P.J.** (1984). A prediction model for parasitic gastro-enteritis in lambs. *International Journal for Parasitology* **14**: 439-445.



- Paterson, S. and Viney, M.E.** (2002). Host immune responses are necessary for density dependence in nematode infections. *Parasitology* **125**: 283-292.
- Pence, D.B.** (1990). Helminth community of mammalian hosts: concepts at the infracommunity, component and compound community levels. In: Esch, G.W., Bush, A.O. and Aho, J.M. (Eds.). *Parasite Communities: Patterns and Processes*. Chapman and Hall, London.
- Penzhorn, B.L.** (2000). Coccidian oocyst and nematode egg counts of free-ranging African buffalo (*Syncerus caffer*) in the Kruger National Park, South Africa. *Journal of the South African Veterinary Association* **71**: 106-108.
- Peters, B.G. and Leiper, J.W.G.** (1940). Variation in dilution-counts of helminth eggs. *Journal of Helminthology* **18**: 117-142.
- Petney, T.N., Van Ark, H. and Spickett, A.M.** (1990). On sampling tick populations: the problem of overdispersion. *Onderstepoort Journal of Veterinary Research* **57**: 123-127.
- Petrov, V.S.** (1976). Invasion of saigas with *Avitellina* and other intestinal cestodes by age and season. *Contributions to the Natural Nidality of Diseases* **8**: 130-131. [In Russian.]
- Petrov, V.S.** (1985). Helminths of saigas and their importance for the epizootology of helminthoses in sheep. Doctoral thesis, Moscow. [In Russian.]
- Pianka, E.R.** (2000). *Evolutionary Ecology*. 6<sup>th</sup> ed. Addison Wesley Longman, San Francisco.
- Plowright, R.C. and Paloheimo, J.E.** (1977). A theoretical study of population dynamics in the sheep tick. *Theoretical Population Biology* **12**: 286-297.
- Poole, J.B.** (1953). Reaction to temperature by infective larvae of *Nematodirus filicollis*, Trichostrongylidae (Nematoda). *Canadian Journal of Comparative Medicine* **20**: 169-172.
- Poulin, R.** (1993). The disparity between observed and uniform distributions: a new look at parasite aggregation. *International Journal for Parasitology* **23**: 937-944.
- Poulin, R.** (1996). Helminth growth in vertebrate hosts: does host sex matter? *International Journal for Parasitology* **26**: 1311-1315.
- Poulin, R.** (1998). *Evolutionary Ecology of Parasites: From Individuals To Communities*. Chapman and Hall, London.
- Priyadko, E.I., Tastanov, T.B. and Baitursunov, K.K.** (1995a). *Helminths of the Saiga and conserved animals of Kazakhstan*. Unpublished report. Institute of Zoology, Almaty, Kazakhstan. [In Russian.]
- Priyadko, E.I., Tastanov, T.B. and Baitursunov, K.K.** (1995b). Possible methods of anthelmintic prophylaxis in saigas. *Proceedings of the National Academy of Sciences of the Republic of Kazakhstan* **3**: 8-11. [In Russian.]
- Pronyaev, A.V., Fandeev, A.A. and Gruzdev, A.R.** (1998). Age related variation of teeth, techniques of age determination. Pp. 209-215 in: Sokolov, V.E. and Zhirnov, L.V. (Eds.). *The Saiga Antelope: Phylogeny, Systematics, Ecology, Conservation and Use*. Russian Academy of Sciences, Moscow. 356pp. [In Russian.]
- Quinnell, R.J., Medley, G.F. and Keymer, A.E.** (1990). The regulation of gastrointestinal helminth populations. *Philosophical Transactions of the Royal Society of London, Series B* **330**: 191-201.

- Quinnell, R.J., Grafen, A. and Woolhouse, M.E.J.** (1995). Changes in parasite aggregation with age: a discrete infection model. *Parasitology* **111**: 635-644.
- Radionov, P.V.** (1973a). On questions of the transmission of helminths between sheep, cattle and saigas in Kustanai *oblast* and natural hosts of some helminths. *Contributions to the Natural Nidality of Diseases* **6**: 121-130. [In Russian.]
- Radionov, P.V.** (1973b). On the transmission of helminths between sheep, cattle and saigas in Kustanai *oblast*. *Transactions of the Kazakhstan Scientific Veterinary Research Institute* **15**: 312-315. [In Russian.]
- Radionov, P.V., Shkunov, P.Ya., Nikitenko, G.I., Sabaev, B.S. and Balkibekov, A.B.** (1968). The epizootology, diagnosis and therapy of nematodiosis in sheep. Pp. 124-126 in: *Proceedings of The Advisory Seminar on the Control of Helminthoses of Domestic Animals*. Kazakhstan Scientific Investigation Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]
- Radostits, O.M., Gay, C.C., Blood, D.C. and Hinchcliff, K.W.** (2000). *Veterinary Medicine: A Textbook of the Diseases of Cattle, Sheep, Pigs, Goats and Horses*. Saunders, London.
- Randolph, S.E.** (2000). Ticks and tick-borne disease systems in space and from space. *Advances in Parasitology* **47**: 217-245.
- Read, A.F., Albon, S.D., Antonovics, J., Apanius, V., Dwyer, G., Holt, R.D., Judson, O., Lively, C.M., Martin-Löf, A., McLean, A.R., Metz, J.A.J., Schmid-Hempel, P., Thrall, P.H., Via, S. and Wilson, K.** (1995). Group report: Genetics and evolution of infectious diseases in natural populations. In: Grenfell, B.T. and Dobson, A.P. (Eds.). *Ecology of Infectious Diseases in Natural Populations*. Cambridge University Press. Pp. 450-477.
- Reinecke, R.K., Snijders, A.J. and Horak, I.G.** (1962). A modification of standard procedures for evaluating the relative efficacy of anthelmintics. *Onderstepoort Journal of Veterinary Research* **29**: 241-257.
- Reinecke, R.K.** (1984). Identification of helminths in ruminants at necropsy. *Journal of the South African Veterinary Association* **55**: 135-143.
- Reinecke, R.K.** (1994). Parasitic control in intensive vs. non-intensive systems – ruminants. *Veterinary Parasitology* **54**: 49-67.
- Renshaw, E.** (1999). Stochastic effects in population models. In: *Advanced Ecological Theory. Principles and Applications*. (Ed. McGlade, J.). Blackwell Science, Oxford. Pp. 64-99.
- Rickard, L.G. and Zimmerman, G.L.** (1992). The epizootology of gastrointestinal nematodes of cattle in selected areas of Oregon. *Veterinary Parasitology* **43**: 271-291.
- Rickard, L.G., Hoberg, E.P., Zimmerman, G.L. and Erno, J.K.** (1987). Late fall transmission of *Nematodirus battus* (Nematoda: Trichostrongyloidea) in western Oregon. *Journal of Parasitology* **73**: 244-247.
- Ringrose, S., Vanderpost, C., Matheson, W.** (1997). Use of image processing and GIS technology to determine the extent and possible causes of land management / fence-line induced degradation problems in the Okavango area, northern Botswana. *International Journal of Remote Sensing* **18**: 2337-2364.

- Roberts, M.G. and Grenfell, B.T.** (1991). The population dynamics of nematode infections of ruminants: periodic perturbations as a model for management. *IMA Journal of Mathematics Applied in Medicine and Biology* **8**: 83-93.
- Roberts, M.G. and Heesterbeek, J.A.P.** (1995). The dynamics of nematode infections of farmed ruminants. *Parasitology* **110**: 493-502.
- Roberts, J.L. and Swan, R.A.** (1982). Quantitative studies of ovine haemonchosis. 3. The interpretation and diagnostic significance of the changes in serial egg counts of *Haemonchus contortus* in a sheep flock. *Veterinary Parasitology* **9**: 211-216.
- Robinson, S.** (2000). Pastoralism and land degradation in Kazakhstan. PhD Thesis, University of Warwick.
- Robinson, S. and Milner-Gulland, E.J.** (2003). Political change and factors limiting numbers of wild and domestic ungulates in Kazakhstan. *Human Ecology* **31**: 87-110.
- Rogers, D.J.** (2000). Satellites, space, time and the African trypanosomiasis. *Advances in Parasitology* **47**: 130-173.
- Roland, J.** (1988). Decline of winter moth populations in North America: direct v indirect effect of introduced parasites. *Journal of Animal Ecology* **57**: 523-531.
- Rose, J.H.** (1963). Observations on the free-living stages of the stomach worm *Haemonchus contortus*. *Parasitology* **53**: 469-481.
- Rose, C.H. and Jacobs, D.E.** (1990). Epidemiology of *Nematodirus* species of sheep in a subarctic climate: development and persistence of larvae on herbage. *Research in Veterinary Science* **48**: 327-330.
- Rose, J.H. and Small, A.J.** (1980). Transmission of *Oesophagostomum* spp. among sows at pasture. *Veterinary Record* **107**: 223-225.
- Rose, J.H. and Small, A.J.** (1985). The distribution of the infective larvae of sheep gastrointestinal nematodes in soil and on herbage and the vertical migration of *Trichostrongylus vitrinus* larvae through the soil. *Journal of Helminthology* **59**: 127-135.
- Rose, C.H., Jacobs, D.E., Jørgensen, R.J. and Nansen, P.** (1984). Studies of helminth parasites of sheep in southern Greenland. *Nordisk Veterinaermedicin* **36**: 77-87.
- Rossanigo, C.E. and Gruner, L.** (1991). Accuracy of two methods for counting eggs of sheep nematode parasites. *Veterinary Parasitology* **39**: 115-121.
- Rousset, F., Thomas, F., De Meeus, T. and Renaud, F.** (1996). Inference of parasite-induced host mortality from distributions of parasite loads. *Ecology* **77**: 2203-2211.
- Rozenzweig, M.L.** (1999). Species diversity. In: *Advanced Ecological Theory. Principles and Applications*. (Ed. McGlade, J.). Blackwell Science, Oxford. Pp. 249-281.
- Rózsa, L., Reiczigel, J. and Majoros, G.** (2000). Quantifying parasites in samples of hosts. *Journal of Parasitology* **86**: 228-232.
- Rubin, R.** (1967). Some observations on the interpretation of fecal egg counts. *American Journal of Veterinary Clinical Pathology* **1**: 145-148.
- Ruzimuradov** (1966). Cited in Irgashev (1973).

**Sadikov, Zh.** (1975). The epizootological situation of helminthoses of sheep in the north-east of Taldy-Kurgan *oblast*. Pp. 290-291 in: *Diseases of Sheep and Means of Their Control in Kazakhstan*. Kazakhstan Scientific Research Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]

**Sage, R.B., Woodburn, M.I.A., Davis, C. and Aebischer, N.J.** (2002). The effect of an experimental infection of the nematode *Heterakis gallinarum* on hand-reared grey partridges *Perdix perdix*. *Parasitology* **124**: 529-535.

**Samson, K.S., Allen, R.W. and Schad, G.A.** (1964). Comparative pathogenicity in *Ovis aries* of homologous and heterologous strains of *Haemonchus* (Nematoda: Trichostrongylidae) from domestic and wild sheep. *The Journal of Parasitology* **50**: 421-426.

**Samuel, W., Pybus, M. and Kocan, A.A.** (Eds.) (2001). *Parasitic Diseases of Wild Mammals* (2<sup>nd</sup> Ed.). Blackwell, Oxford.

**Satubaldin, Kh.** (1975). On the study of helminths and helminthoses of sheep in holdings in East Kazakhstan *oblast*. Pp. 291-292 in: *Diseases of Sheep and Means of Their Control in Kazakhstan*. Kazakhstan Scientific Research Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]

**Saunders, L.M., Tompkins, D.M. and Hudson, P.J.** (2001). Strategies for nematode transmission: selective migration of *Trichostrongylus tenuis* infective larvae. *Journal of Helminthology* **75**: 367-372.

**Scholl, V.A., Osipov, P.P. and Zhevnerov, V.V.** (1979). Natural hosts of helminthoses of ruminants on Barsa-Kel'mes island. *Contributions to the Natural Nidality of Diseases* **10**: 161-170. [In Russian.]

**Scott, M.E.** (1987). Regulation of mouse colony abundance by *Heligmosomoides polygyrus*. *Parasitology* **95**: 111-124.

**Shaikenov, B.** (1975). On the epizootology of nematodiosis of sheep in the foothill zone of the Dzhungarian Alatau. Pp. 324-326 in: *Diseases of Sheep and Means of Their Control in Kazakhstan*. Kazakhstan Scientific Research Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]

**Shaikenov, B., Milner-Gulland, E.J., Torgerson, P, Morgan, E. and Bekturganov, B.** (1999). Transmission of parasites between the Betpak-Dala population of saigas and domestic animals. Pp. 194-195 in: *Proceedings of the Conference on Problems and Sustainable Use of the Animal Fauna of Kazakhstan*. Almaty, 6<sup>th</sup> April 1999. Institute of Zoology, Almaty, Kazakhstan. [In Russian.]

**Shaikenov, B.S., Vaganov, T.F. and Torgerson, P.R.** (1999). Cystic echinococcosis in Kazakhstan: An emerging disease since independence from the Soviet Union. *Parasitology Today* **15**: 172-174.

**Sharkhuu, T.** (2001). Helminths of goats in Mongolia. *Veterinary Parasitology* **101**: 161-169.

**Shaw, D.J. and Dobson, A.P.** (1995). Patterns of macroparasite abundance and aggregation in wildlife populations: a quantitative review. *Parasitology* **111**: S111-S133.

**Shaw, D.J., Grenfell, B.T. and Dobson, A.P.** (1998). Patterns of macroparasite aggregation in wildlife host populations. *Parasitology* **117**: 597-610.

- Shaw, J.L. and Moss, R.** (1989). The role of parasite fecundity and longevity in the success of *Trichostrongylus tenuis* in low density red grouse populations. *Parasitology* **99**: 252-258.
- Shaw, J.L., Moss, R. and Pike, A.W.** (1989). Development and survival of the free-living stages of *Trichostrongylus tenuis*, a caecal parasite of red grouse *Lagopus lagopus scoticus*. *Parasitology* **99**: 105-113.
- Shorb, D.A.** (1943). Survival on grass plots of eggs and preinfective larvae of the common sheep stomach worm, *Haemonchus contortus*. *Journal of Parasitology* **29**: 284-289.
- Siegel, S. and Castellan, N.J.** (1988). *Non-parametric Statistics for the Behavioural Sciences*. (2<sup>nd</sup> Ed.) McGraw-Hill, New York.
- Silangwa, S.M. and Todd, A.C.** (1964). Vertical migration of trichostrongylid larvae on grasses. *The Journal of Parasitology* **50**: 278-285.
- Silangwa, S.M. and Todd, A.C.** (1968). Vertical migration of trichostrongylid larvae on grasses. *The Journal of Parasitology* **50**: 278-285.
- Silverman, P.H. and Campbell, J.A.** (1959). Studies on parasitic worms of sheep in Scotland. I. Embryonic and larval development of *Haemonchus contortus* at constant conditions. *Parasitology* **49**: 23-38.
- Sinclair, A.R.E.** (1989). Population regulation in animals. In: Cherrett, J.M. (Ed.). *Ecological Concepts. The Contribution of Ecology to an Understanding of the Natural World*. Blackwell Scientific Publications, Oxford.
- Singh, V., Kumar, S. and Yadav, C.L.** (1993). Establishment of camel isolates of *Haemonchus longistipes* and *Trichostrongylus colubriformis* in goats. *Veterinary Parasitology* **51**: 69-75.
- Skira, V.N.** (1995). Population dynamics of *Nematodirus spathiger* in lambs. *Information Bulletin Kharkov 1994*. Experimental Institute of Clinical Veterinary Medicine, Ukrainian Academy of Sciences. [In Russian.]
- Skrjabin, K.I., Shikobalova, N.P. and Schulz, R.S.** (1954). *Essentials of Nematodology. Vol. III. Trichostrongylids of Animals and of Man*. USSR Academy of Sciences Press, Moscow. [In Russian.]
- Skrjabin, K.I., Shikobalova, N.P. and Schulz, R.S.** (1960). *Essentials of Nematodology. Vol. III. Trichostrongylids of Animals and of Man*. USSR Academy of Sciences Press, Moscow. English Version: The Israel Program for Scientific Translations, Jerusalem.
- Slater, A.F.G. and Keymer, A.E.** (1988). The influence of protein deficiency on immunity to *Heligmosomoides polygyrus* (Nematoda) in mice. *Parasite Immunology* **10**: 507-522.
- Slocombe, J.O.D.** (1974). Overwintering of bovine gastrointestinal nematodes in southwestern Ontario. *Canadian Journal of Comparative Medicine* **38**: 90-93.
- Sloss, M.W., Kemp, R.L. and Zajac, A.M.** (1994). *Veterinary Clinical Parasitology*. 6<sup>th</sup> Ed. American Association of Veterinary Parasitology. Iowa State University Press, Ames. 198 pp.
- Sludskii, A.A.** (1955). The Saiga in Kazakhstan. *Proceedings of the Zoological Institute of the Kazakhstan Academy of Sciences* **4**: 18-55. Alma-Ata, USSR. [In Russian]

- Sludskii, A.A.** (1962). The relationship between predator and prey. *Proceedings of the Zoological Institute of the Kazakhstan Academy of Sciences* **7**: 24-143. (In Russian; cited in Bekenov *et al*, 1998.)
- Smeal, M.G., Hotson, I.K., Mylrea, P.J., Jackson, A.R., Campbell, N.J. and Kirton, H.C.** (1977). Studies on nematode infections of beef cattle in New South Wales. *Australian Veterinary Journal* **53**: 566-573.
- Smith, G.** (1988). The population biology of the parasitic stages of *Haemonchus contortus*. *Parasitology* **96**: 185-195.
- Smith, G.** (1989). The population biology of the parasitic stages of *Ostertagia circumcincta*. *International Journal for Parasitology* **19**: 385-393.
- Smith, G.** (1990). The population biology of the free-living phase of *Haemonchus contortus*. *Parasitology* **101**: 309-316.
- Smith, G.** (1994). Population biology of the parasitic phase of trichostrongylid nematode parasites of cattle and sheep. *International Journal for Parasitology* **24**: 167-178.
- Smith, G. and Galligan, D.T.** (1988). Mathematical models of the population biology of *Ostertagia ostertagi* and *Teladorsagia circumcincta*, and the economic evaluation of disease control strategies. *Veterinary Parasitology* **27**: 73-83.
- Smith, G. and Grenfell, B.T.** (1994). Modelling of parasite populations: gastrointestinal nematode models. *Veterinary Parasitology* **54**: 127-143.
- Smith, G. and Guerrero, J.** (1993). Mathematical models of the population biology of *Ostertagia ostertagi* and the significance of aggregated parasite distributions. *Veterinary Parasitology* **46**: 243-257.
- Smith, G. and Scott, M.E.** (1994). Model behavior and the basic reproduction ratio. In Scott, M.E. and Smith, G. (Eds.) *Parasitic and Infectious Diseases: Epidemiology and Ecology*. Academic Press, San Diego. Pp. 21-28.
- Smith, G., Grenfell, B.T. and Anderson, R.M.** (1986). The development and mortality of the non-infective free-living stages of *Ostertagia ostertagi* in the field and in laboratory culture. *Parasitology* **92**: 471-482.
- Smith, G., Grenfell, B.T. and Anderson, R.M.** (1987). The regulation of *Ostertagia ostertagi* in calves: density-dependent control of fecundity. *Parasitology* **95**: 373-388.
- Smith, H.J. and Archibald, R.McG** (1965). The overwinter survival of ovine gastro-intestinal parasites in the maritime provinces. *Canadian Veterinary Journal* **6**: 257-267.
- Sokal, R.R. and Rohlf, F.J.** (1995). *Biometry: The Principles and Practice of Statistics in Biological Research*. 3<sup>rd</sup> Ed. Freeman, New York. 887pp.
- Sokolov, V.E. and Zhirnov, L.V.** (Eds.) (1998). *The Saiga Antelope. Phylogeny, Systematics, Conservation and Land Use*. Moscow. [In Russian]
- Soulsby, E.J.L.** (1982). *Helminths, Arthropods and Protozoa of Domesticated Animals*. Balliere Tindall, London.

- Southcott, W.H. and Barger, I.A.** (1975). Control of nematode parasites by grazing management. II. Decontamination of cattle pastures by varying periods of grazing with the alternative host. *International Journal for Parasitology* **5**: 39-44.
- Sreter, T., Monlar, V. and Kassai, T.** (1994). The distribution of nematode egg counts and larval counts in grazing sheep and their implications for parasite control. *International Journal for Parasitology* **24**: 103-108.
- Stampa, S. and Linde, S.** (1972). A contribution towards the diagnosis of trichostrongylosis in flocks of sheep under natural conditions. *Journal of the South African Veterinary Association* **43**: 95-99.
- Stear, M.J., Bishop, S.C., Duncan, J.L., McKellar, Q.A. and Murray, M.** (1995a). The repeatability of faecal egg counts, peripheral eosinophil counts, and plasma pepsinogen concentrations during deliberate infections with *Ostertagia circumcincta*. *International Journal for Parasitology* **25**: 375-380.
- Stear, M.J., Bairden, K., Duncan, J.L., Gettinby, G., McKellar, Q.A., Murray, M. and Wallace, D.S.** (1995b). The distribution of faecal nematode egg counts in Scottish Blackface lambs following natural, predominantly *Ostertagia ostertagi* infection. *Parasitology* **110**: 573-581.
- Stear, M.J., Bairden, K., Duncan, J.L., Holmes, P.H., McKellar, Q.A., Park, M., Strain, S., Murray, M., Bishop, S.C. and Gettinby, G.** (1997). How hosts control worms. *Nature* **389**: 27.
- Stear, M.J. and Bishop, S.C.** (1999). The curvilinear relationship between worm length and fecundity of *Teladorsagia circumcincta*. *International Journal for Parasitology* **29**: 777-780.
- Sterud, E., Mo, T.A., Collins, C.M. and Cunningham, C.O.** (2002). The use of host specificity, pathogenicity, and molecular markers to differentiate between *Gyrodactylus salaries* Malmberg, 1957 and *G. thymalli* Zitnan, 1960 (Monogenea: Gyrodactylidae). *Parasitology* **124**: 203-213.
- Stoll, N.R.** (1930). On methods of counting nematode ova in sheep dung. *Parasitology* **22**: 116-136.
- Stromberg, B.E.** (1997). Environmental factors influencing transmission. *Veterinary Parasitology* **72**: 247-264.
- Stromberg, B.E. and Corwin, R.M.** (1993). Epizootiology of *Ostertagia ostertagi* in cow-calf production systems in the American Midwest. *Veterinary Parasitology* **46**: 297-302.
- Stroud, R.K.** (1978). Parasites and associated pathology in pinnipeds stranded along the Oregon coast. *Journal of Wildlife Diseases* **14**: 292-298.
- Suarez, V.H. and Buseti, M.R.** (1995). The epidemiology of helminth infections of growing sheep in Argentina's western pampas. *International Journal for Parasitology* **25**: 489-494.
- Suarez, V.H., Cabaret, J. and Gruner, L.** (1995). Morphological polymorphism in the nematode *Teladorsagia circumcincta* in relation to age of larvae, infection mode and lamb characteristics in experimental conditions. *International Journal for Parasitology* **25**: 1173-1177.

- Sukhdeo, M.V.K. and Bansemir, A.D.** (1996). Critical resources that influence habitat selection decisions by gastrointestinal helminth parasites. *International Journal for Parasitology* **26**: 483-498.
- Sutmoller, P., Thomson, G.R., Hargreaves, S.,K., Foogin, C.M. and Anderson, E.C.** (2000). The foot-and-mouth disease risk posed by African buffalo within wildlife conservancies to the cattle industry of Zimbabwe. *Preventive Veterinary Medicine* **44**: 43-60.
- Swan, R.A.** (1970). An improved method for the collection of faecal samples from sheep. *Australian Veterinary Journal* **46**: 25-26.
- Sykes, A.R.** (1978). The effect of subclinical parasitism in sheep. *The Veterinary Record* **102**: 32-34.
- Sykes, A.R.** (1987). Endoparasites and herbivore nutrition. In: *The Nutrition of Herbivores* (Eds. Hacker, J.B. and Ternouth, J.H.). Academic Press, Australia. Pp. 211-232.
- Szanto, J., Mohan, L.V.P. and Levine, N.D.** (1964). Prevalence of coccidian and gastrointestinal nematodes in beef cattle in Illinois and their relation to shipping fever. *Journal of the American Veterinary Medical Association* **144**: 741-746.
- Tallis, G.M. and Leyton, M.K.** (1966). A stochastic approach to the study of parasite populations. *Journal of Theoretical Biology* **13**: 251-260.
- Tarazona, J.M.** (1986). A method for the interpretation of parasite egg counts in faeces of sheep. *Veterinary Parasitology* **22**: 113-119.
- Taylor, E.L.** (1939). Technique for the estimation of pasture infestation by strongyloid larvae. *Parasitology* **31**: 473-478.
- Taylor, L.R.** (1961). Aggregation, variance and the mean. *Nature* **189**: 732-735.
- Taylor, L.R., Woiod, I.P. and Perry, J.N.** (1978). The density-dependence of spatial behaviour and the rarity of randomness. *Journal of Animal Ecology* **47**: 383-406.
- Taylor, E.L., Catchpole, J., Marshall, R., Norton, C.C. and Green, J.** (1995). *Eimeria* spp. of sheep. Pp. 25-39 in Eckert, J., Taylor, M., Braun, R., Shirley, M.W. and Coudert, P. (Eds.). *Guidelines in Techniques in Coccidiosis Research*. European Commission, Brussels.
- Tembely, S., Lahlou-kassi, A., Rege, J.E.O., Sovani, S., Diedhiou, M.L. and Baker, R.L.** (1997). The epidemiology of nematode infections in sheep in a cool tropical environment. *Veterinary Parasitology* **70**: 129-141.
- Tembely, S., Lahlou-kassi, A., Rege, J.E.O., Mukasa-Mugerwa, E., Anindo, D., Sovani, S. and Baker, R.L.** (1998). Breed and season effects on the peri-parturient rise in nematode egg output in indigenous ewes in a cool tropical environment. *Veterinary Parasitology* **77**: 123-132.
- Thamsborg, S.M., Jørgensen, R.J., Waller, P.J. and Nansen, P.** (1996). The influence of stocking rate on gastrointestinal nematode infections of sheep over a 2-year grazing period. *Veterinary Parasitology* **67**: 207-224.
- Tharaldsen, J.** (1976). The epidemiology of trichostrongylid infections in young cattle in Norway. *Acta Veterinaria Scandinavica (Supplement)* **61**: 1-21.



- Tharaldsen, J. and Helle, O.** (1984). Epidemiological investigations of trichostrongylid infections in young cattle in different parts of Norway. *Acta Veterinaria Scandinavica* **25**: 164-186.
- Thienpont, D., Rochette, F. and Vanparijs, O.F.J.** (1979). *Diagnosing Helminthiasis by Coprological Examination*. Janssen Research Foundation, Beerse, Belgium.
- Thomas, R.J.** (1979). A comparative study of the life histories of *Nematodirus battus* and *N. filicollis*, nematode parasites of sheep. *Parasitology* **49**: 374-386.
- Thomas, R.J. and Boag, B.** (1973). Epidemiological studies on gastro-intestinal nematode parasites of sheep. The control of infection in lambs on contaminated pasture. *Research in Veterinary Science* **15**: 238-249.
- Thomas, R.J. and Stevens, A.J.** (1960). Ecological studies on the development of the pasture stages of *Nematodirus battus* and *N. filicollis*, nematode parasites of sheep. *Parasitology* **50**: 31-49.
- Thomas, R.T.** (1974). The role of climate in the epidemiology of nematode parasites of ruminants. In: Taylor, A.E.R. and Muller, R. (Eds.). *The effects of meteorological factors upon parasites. Symposia of the British Society for Parasitology, Volume 12*. Blackwell Scientific Publications, Oxford.
- Thomsen, L.E., Mejer, H., Wendt, S., Roepstorff, A. and Hindsbo, O.** (2001). The influence of helminth parasites in pigs on permanent pasture during two consecutive summers. *Veterinary Parasitology* **99**: 129-146.
- Thomson, G.R.** (1995). Overview of foot and mouth disease in southern Africa. *Revue Scientifique et Technique de L'Office International des Epizooties* **14**: 503-520.
- Thorne, E.T. and Williams, E.S.** (1988). Disease and endangered species: The black-footed ferret as a recent example. *Conservation Biology* **2**: 66-73.
- Thrusfield, M.** (1995). *Veterinary Epidemiology*. (2<sup>nd</sup> Ed.) Blackwell, Oxford. 483pp.
- Tompkins, D.M. and Begon, M.** (1999). Parasites can regulate wildlife populations. *Parasitology Today* **15**: 311-313.
- Tompkins, D.M., Greenman, J.V. and Hudson, P.J.** (2001). Differential impact of a shared nematode parasite on two gamebird hosts: implications for apparent competition. *Parasitology* **122**: 187-193.
- Tompkins, D.M., Dobson, A.P., Arneberg, P., Begon, M.E., Cattadori, I.M., Greenman, J.V., Heesterbeek, J.A.P., Hudson, P.J., Newborn, D., Pugliese, A., Rizzoli, A.P., Rosà, R., Rosso, F. and Wilson, K.** (2002a). Parasites and host population dynamics. Pp. 45-62 in: Hudson, P.J., Rizzoli, A., Grenfell, B.T., Heesterbeek, H. and Dobson, A.P. (Eds.). *The Ecology of Wildlife Diseases*. Oxford University Press.
- Tompkins, D.M., Sainsbury, A.W., Nettleton, P., Buxton, D. and Gurnell, J.** (2002b). Parapoxvirus causes deleterious disease in red squirrels associated with UK population declines. *Proceedings of the Royal Society of London Series B: Biological Sciences* **269**: 529-533.
- Torgerson, P.R., Pilkington, J., Gulland, F.M.D. and Gemmell, M.A.** (1995). Further evidence for the long distance dispersal of taeniid eggs. *International Journal for Parasitology* **25**: 265-267.

- Torgerson, P.R., Shaikenov, B.S., Baitursinov, K.K. and Abdybekova, A.M.** (2002). The emerging epidemic of echinococcosis in Kazakhstan. *Transactions of the Royal Society of Tropical Medicine & Hygiene* **96**: 124-128.
- Torgerson, P.R., Shaikenov, B.S., Rysmukhambetova, A.T., Ussenbayev, A.E., Abdybekova, A.M., and Burtisunov, K.K.** (2003a). Modelling the transmission dynamics of *Echinococcus granulosus* in dogs in Kazakhstan. *Parasitology* **126**: 417-424.
- Torgerson, P.R., Burtisunov, K.K., Shaikenov, B.S., Rysmukhambetova, A.T., Abdybekova, A.M. and Ussenbayev, A.E.** (2003b). Modelling the transmission dynamics of *Echinococcus granulosus* in sheep and cattle in Kazakhstan. *Veterinary Parasitology* **114**: 143-153.
- Trach, V.N.** (1964). Data cited in Trach (1983).
- Trach, V.N.** (1983). *Parasitic Strongylate Larvae of Domestic Ruminants*. Ukrainian Academy of Sciences, I.I. Shmal'gauzena Institute of Zoology, Kiev, USSR. [In Russian.]
- Trach, V.N.** (1986). *Ecologo-faunal Characteristics of Sexually Mature Strongylates of Domestic Ruminants of the Ukraine*. Ukrainian Academy of Sciences, I.I. Shmal'gauzena Institute of Zoology, Kiev, USSR. [In Russian.]
- Urquhart, G.M., Armour, J., Duncan, L., Dunn, A.M. and Jennings, F.W.** (1996). *Veterinary Parasitology* (2<sup>nd</sup> Ed.). Longman, Avon.
- Valderrábano, J., Delfa, R. and Uriarte, J.** (2002). Effect of feed intake on the development of gastrointestinal parasitism in growing lambs. *Veterinary Parasitology* **104**: 327-338.
- Van Houtert, M.F.J.** (1997). Effects of diet on gastrointestinal nematode infection in ruminants. In: Barrell (Ed.). *Sustainable Control of Internal Parasites in Ruminants*. Lincoln University, New Zealand. Pp. 183-201.
- Van Houtert, M.F.J. and Sykes, A.R.** (1996). Implications of nutrition for the ability of ruminants to withstand gastrointestinal nematode infections. *International Journal for Parasitology* **26**: 1151-1168.
- Van Schalkwyk, P.C., Geysler, T.L., Davies, P.V.A. and Recio, M.** (1981). The efficacy of anthelmintics against *Thysaniezia giardi* in South Africa. *Journal of the South African Veterinary Association*, Sept. 1981, 207-209.
- Van Wyk, J.A., Coles, G.C. and Kreczek, R.C.T.** (2002). Can we slow the development of anthelmintic resistance? An electronic debate. *Trends in Parasitology* **18**: 336-337.
- Vercruyse, J. and Claerebout, E.** (1997). Immunity development against *Ostertagia ostertagi* and other gastrointestinal nematodes in cattle. *Veterinary Parasitology* **72**: 309-326.
- Vercruyse, J. Hilderson, H. and Claerebout, E.** (1995). Effect of chemoprophylaxis with avermectins on the immune response to gastrointestinal nematodes in first-season grazing calves. *Veterinary Parasitology* **58**: 35-48.
- Viljoen, J.H.** (1972). Morphology of the free-living stages of *Nematodirus spathiger* with some observations on their development under laboratory conditions. *Journal of the South African Veterinary Association* **43**: 87-94.

**Viney, M. and Read, A.** (2003). Parasite variation: immunological and ecological significance. *Parasitology (Suppl.)* **125**: S1-S2.

**Vlassoff, A. and Bisset, S.A.** (1991). Basic principles of parasite epidemiology. In: Petersen, G.V. (ed.) Practical Aspects of Parasite Control. *Proceedings of the 21<sup>st</sup> Sheep and Beef Cattle Seminar, Lincoln University, New Zealand, July 1991*. Publication no. 134, Veterinary Continuing Education. Massey University, New Zealand.

**Vlassoff, A., Bisset, S.A. and McMurty, L.W.** (1999). Faecal egg counts in Angora goats following natural or experimental challenge with nematode parasites: within-flock variability and repeatabilities. *Veterinary Parasitology* **84**: 113-123.

**Waid, D.D., Pence, D.B. and Warren, R.J.** (1985). Effects of season and physical condition on the gastrointestinal helminth community of white-tailed deer from the Texas Edwards plateau. *Journal of Wildlife Diseases* **21**: 264-273.

**Wakelin, D.** (1994). Host populations: genetics and immunity. In: *Parasitic and Infectious Diseases: Epidemiology and Ecology* (Eds. Scott, M.E. and Smith, G.). Academic Press, San Diego. Pp. 83-100.

**Wallace, D.S., Bairden, K., Duncan, J.L., Fishwick, G., Holmes, P.H., McKellar, Q.A., Murray, M., Parkins, J.J. and Stear, M.J.** (1995). Influence of supplementation with dietary soybean meal on resistance to haemonchosis in Hampshire down lambs. *Research in Veterinary Science* **58**: 232-237.

**Waller, P.J., Dobson, R.J., Donald, A.D. and Thomas, R.J.** (1981). Populations of strongyloid nematode infective stages in sheep pastures: comparison between direct pasture sampling and tracer lambs as estimators of larval abundance. *International Journal for Parasitology* **11**: 359-367.

**Waller, P.J. and Donald, A.D.** (1970). Egg size and desiccation survival in *Trichostrongylus colubriformis* (Nematoda: Trichostrongylidae). *Parasitology* **61**: 205-209.

**Waruiru, R.M., Thamsborg, S.M., Nansen, P., Kyvsgaard, N.C., Bogh, H.O., Munyua, W.K. and Gathuma, J.M.** (2001). The epidemiology of gastrointestinal nematodes of dairy cattle in central Kenya. *Tropical Animal Health and Production* **33**: 173-187.

**Welsh, A.H., Cunningham, R.B., Donnelly, C.F. and Lindenmayer, D.B.** (1996). Modelling the abundance of rare species: statistical models for counts with extra zeros. *Ecological Modelling* **88**: 297-308.

**Williams, J.C.** (1997). Anthelmintic treatment strategies: current status and future. *Veterinary Parasitology* **72**: 461-477.

**Williams, J.C. and Bilkovich, F.R.** (1973). Distribution of *Ostertagia ostertagi* infective larvae on pasture herbage. *American Journal of Veterinary Research* **34**: 1337-1344.

**Williams, B.G. and Dye, C.** (1994). Maximum likelihood for parasitologists. *Parasitology Today* **10**: 489-493.

**Williams, E.S., Thorne, E.T., Appel, M.J.G. and Belitsky, D.W.** (1988). Canine distemper in black-footed ferrets (*Mustela nigripes*) from Wyoming. *Journal of Wildlife Diseases* **24**: 385-398.

- Williams, E.S., Anderson, S.L., Cavender, J., Lynn, C., Hearn, C. and Appel, M.J.G.** (1996). Vaccination of black-footed ferrets (*Mustela nigripes*) X Siberian polecat (*M. eversmanni*) hybrids and domestic ferrets (*M. putorius furo*) against canine distemper. *Journal of Wildlife Diseases* **32**: 417-423.
- Wilson, G.I.** (1957). Comparison of the DCF and McMaster egg-counting techniques as applied to nematode parasitism of sheep. *The Journal of Parasitology* **43** (Suppl.): 20.
- Wilson, K., Grenfell, B.T. and Shaw, D.J.** (1996). Analysis of aggregated parasite distributions: a comparison of methods. *Functional Ecology* **10**: 592-601.
- Wilson, K. and Grenfell, B.T.** (1997). Generalized linear modelling for parasitologists. *Parasitology Today* **13**: 33-38.
- Wilson, K., Bjørnstad, O.N., Dobson, A.P., Merler, S., Poglayen, G., Randolph, S.E., Read, A.F. and Skorpning, A.** (2002). Heterogeneities in macroparasite infections: patterns and processes. In: Hudson, P.J., Rizzoli, A., Grenfell, B.T., Heesterbeek, H. and Dobson, A.P. (Eds.). *The Ecology of Wildlife Diseases*. Oxford University Press.
- Wolfe, A., Hogan, S., Maguire, D., Fitzpatrick, C., Vaughan, L., Wall, D., Hayden, T.J. and Mulcahy, G.** (2001). Red foxes (*Vulpes vulpes*) in Ireland as hosts for parasites of potential zoonotic and veterinary significance. *Veterinary Record* **149**: 759-763.
- Woolhouse, M.E.J.** (1992). A theoretical framework for the immunoepidemiology of helminth infection. *Parasite Immunology* **14**: 563-578.
- Yacob, H.T., Duranton-Grisez, C., Prevot, F., Bergeaud, J.P., Bleuart, C., Jacquiet, Ph., Dorchie, Ph. and Hoste, H.** (2002). Experimental concurrent infection of sheep with *Oestrus ovis* and *Trichostrongylus colubriformis*: negative interactions between parasite populations and related changes in the cellular responses of nasal and digestive mucosae. *Veterinary Parasitology* **104**: 307-317.
- Young, R.R., Nicholson, R.M., Tweedie, R.L. and Schuh, H.J.** (1980a). Quantitative modelling and prediction of development times of the free-living stages of *Ostertagia ostertagi* under controlled and field conditions. *Parasitology* **81**: 493-505.
- Young, R.R., Anderson, N., Overend, D., Tweedie, R.L., Malafant, K.W.J. and Preston, G.A.N.** (1980b). The effect of temperature on times to hatching of eggs of the nematode *Ostertagia circumcincta*. *Parasitology* **81**: 477-49.
- Zarlenga, D.S., Hoberg, E.P., Stringfellow, F. and Lichtenfels, J.R.** (1998). Comparisons of two polymorphic species of *Ostertagia* and phylogenetic relationships within the Ostertagiinae (Nematoda: Trichostrongyloidea) inferred from ribosomal DNA repeat and mitochondrial DNA sequences. *The Journal of Parasitology* **84**: 806-812.
- Zarlenga, D.S., Chute, M.B., Gasbarre, L.C. and Boyd, P.C.** (2001). A multiplex PCR assay for differentiating economically important gastrointestinal nematodes of cattle. *Veterinary Parasitology* **97**: 199-209.
- Zhirnov, L.V.** (1982). *Restored to Life*. Moscow. [In Russian]
- Zhumadilov, K.** (1975). On the epizootology of important helminthoses of sheep in Alma-Ata oblast. Pp. 169-170 in: *Diseases of Sheep and Means of Their Control in Kazakhstan*. Kazakhstan Scientific Research Veterinary Institute, Ministry of Agriculture of the Republic of Kazakhstan, Almaty. [In Russian.]

**Ziborov, N.A.** (1996). The distribution, seasonal and age dynamics of associative disease caused by *Eimeria*, nematodes and cestodes of the gastrointestinal tract of sheep in south Kazakhstan. Pp.13-23 in: *Invasive diseases of Domestic Animals and Their Prophylaxis. A Collection of Scientific Transactions*. Kazakhstan Scientific Investigation Veterinary Institute, Kazakhstan Academy of Agricultural Science, Almaty. [In Russian.]

**Zimmerman, G.L., Worley, D.E., Rickard, L.G.** (1993). Epidemiology of *Ostertagia* in the Northwestern USA. *Veterinary Parasitology* **46**: 303-311.